

A SELECTION PRINCIPLE FOR WEAK KAM SOLUTIONS VIA FREIDLIN–WENTZELL LARGE DEVIATION PRINCIPLE OF INVARIANT MEASURES*

YUAN GAO[†] AND JIAN-GUO LIU[‡]

Abstract. This paper reinterprets the Freidlin–Wentzell variational construction of the rate function in the large deviation principle for invariant measures from the weak KAM perspective. Through a one-dimensional irreversible diffusion process on a torus, we explicitly characterize essential concepts in the weak KAM theory, such as the Peierls barrier and the projected Mather/Aubry/Mañé sets. The weak KAM representation of the Freidlin–Wentzell variational construction of the rate function is discussed based on the global adjustment for the boundary data and the local trimming from the lifted Peierls barriers. This rate function gives the maximal Lipschitz continuous viscosity solution to the corresponding stationary Hamilton–Jacobi equation (HJE), satisfying the Freidlin–Wentzell variational formula for the boundary data on the Aubry set. Choosing meaningful self-consistent boundary data at each local attractor is essential to select a unique weak KAM solution to stationary HJE. This selected viscosity solution also serves as the global energy landscape of the original stochastic process. This selection for stationary HJEs can be described by first taking the long time limit and then taking the zero noise limit, which also provides a special construction of vanishing viscosity approximation.

Key words. Aubry–Mather theory, maximal Lipschitz continuous viscosity solution, global energy landscape, discrete weak KAM problem, irreversible process, Lax–Oleinik semigroup

MSC codes. 37J50, 60F10, 49L25

DOI. 10.1137/22M1519717

1. Introduction. The classical Kolmogorov–Arnold–Moser (KAM) theorem studied the existence of quasi-periodic solutions of a perturbed integral system. A canonical map converting original phase variables (p, x) to action-angle variables (P, X) can be used to transform a perturbed Hamiltonian dynamics into a (nearly) integrable system in terms of the action-angle variable. This is a classical way to study the perturbed Hamiltonian dynamics, pioneered by Kolmogorov, Arnold, and Moser. The canonical map is globally implicitly solved from a generating function $u(P, x)$, which solves an associated stationary Hamilton–Jacobi equation (HJE) for each action variable P in the classical sense [Arn13, Eva08]. This procedure is, in general, very hard and can only be taken in a small perturbation way. For general Hamiltonian systems far away from an integrable one, the Aubry–Mather theory developed by Aubry [Aub83] and Mather [Mat82] introduced various action minimizing sets and effective Hamiltonian $\bar{H}(P)$ for the corresponding Lagrangian dynamics to obtain a global understanding of the general Hamiltonian dynamics. Instead of finding the classical solution to stationary HJEs with an effective Hamiltonian $\bar{H}(P)$, the notion of a global nondifferential solution defined in the viscosity sense was introduced by

*Received by the editors September 2, 2022; accepted for publication (in revised form) June 13, 2023; published electronically November 2, 2023.

<https://doi.org/10.1137/22M1519717>

Funding: The work of the first author was supported by NSF under award DMS-2204288. The work of the second author was supported by NSF under award DMS-2106988.

[†]Department of Mathematics, Purdue University, West Lafayette, IN 47907 USA (gao662@purdue.edu).

[‡]Department of Mathematics and Department of Physics, Duke University, Durham, NC 22708 USA (jliu@math.duke.edu).

Crandall and Lions [CL83]. Solving the family of stationary HJEs with an effective Hamiltonian in the viscosity sense has many important applications, for instance, the cell problem for the homogenization theory by Lions, Papanicolaou, and Varadhan [LPV86] in the late 1980s.

The above celebrated results on stationary HJEs lead to the development of the weak KAM theory, pioneered particularly by Fathi [Fat97, Fat98], Mañé [Man96], and E [E99]. It is well known that solutions to dynamic HJEs can be represented in terms of the Lax–Oleinik variational formula, which computes the least action of the corresponding Lagrangian at a finite time. The weak KAM theorem by Fathi [Fat98, Fat08] proved convergence from the Lax–Oleinik semigroup representation for the dynamic HJE to a variational representation of solutions to the stationary HJE, known as weak KAM solutions. In other words, weak KAM solutions are invariant solutions for the Lax–Oleinik semigroup; see (4.18). This variational representation for stationary solutions to HJE uses the Mañé potential (3.4) to compute the least action path in an undefined time horizon. Particularly, if one solves the least action problem (backward characteristic of the Hamiltonian dynamics) as $t \rightarrow -\infty$, which tracks back to some invariant sets of the Hamiltonian dynamics, then the Mañé potential becomes the so-called Peierls barrier with an infinite time horizon. More importantly, those backward characteristics obtained through PDE methods can be used to characterize invariant sets in the Aubry–Mather theory for the original dynamical system.

Finding the integrable structure and characterizing those invariant sets of the original dynamics via the variational representation of the globally defined stationary solution to HJE is the central idea in the development of the weak KAM theory and thus is viewed as a generalization of KAM theory in terms of the “Hamilton–Jacobi methods.” By using the concept of the projected Aubry set, the stationary variational representation only relies on the boundary values of the solution $W(x_j)$ on the projected Aubry set \mathcal{A} and the Peierls barriers

$$(1.1) \quad W(x) = \min_{x_j \in \text{invariant states}} W(x_j) + h(x; x_j).$$

This variational representation formula (1.1) was indeed already derived in 1969 in the Freidlin–Wentzell theory for the large deviation principle in the zero noise limit of the invariant measures for diffusion processes [VF69, VF70]. The Freidlin–Wentzell theory comprehensively studied the global quasi-potentials that were globally defined and were later called the Mañé potentials during the development of the weak KAM theory in the late 1990s. The local quasi-potentials within the basin of attraction of each stable state are widely used in computing the barriers for exit problems of a stochastic dynamics. Using quasi-potentials for each basin of attraction of stable states, the Peierls barrier can be computed and can be used to construct the rate function $W(x)$ for the large deviation principle of invariant measures for those stochastic processes; cf. [FW12, Chapter 6, Theorem 4.3].

This paper focuses on reinterpreting the Freidlin–Wentzell variational construction of the rate function for invariant measures in the large deviation principle from the weak KAM perspective. Through a simple one-dimensional irreversible diffusion process on a torus, we explicitly characterize all the essential concepts in the weak KAM theory, such as the projected Aubry/Mather sets, the variational representation, and the unique selection principle for boundary data on the projected Aubry set provided by the large deviation principle. These weak KAM characterizations also, in turn, help us understand the global properties of the rate function in the large deviation principle through a geometric/dynamic viewpoint and the construction of

the global energy landscape, which guides the most probable path/states in the zero noise limit of the stochastic process.

We first clarify that to study the rate function for the large deviation principle, we are only interested in the critical energy level, i.e., the critical Mañé value $c^* = 0$; see section 3.1.1. Then the stationary HJE is

$$(1.2) \quad H(W'(x), x) = W'(W' - U) = 0.$$

Here the Hamiltonian $H(p, x) = p(p - U'(x))$ can be derived from the WKB expansion for the family of invariant measures $\pi_\varepsilon(x) = e^{-\frac{W_\varepsilon(x)}{\varepsilon}}$. The corresponding Lagrangian is also called the Mañé Lagrangian,

$$(1.3) \quad L(s, x) = \frac{1}{4}(s + U'(x))^2.$$

The most distinguishing feature is that the Mañé Lagrangian $L(s, x) \geq 0$ and $L = 0$ if and only if $s = -U'(x)$. This reduces the action minimizing path to a first-order ODE problem. Although it is not directly related to the large deviation principle, there are also other dynamics corresponding to the effective Hamiltonian $\bar{H}(P) > c^* = 0$. This defines different invariant sets and action minimizing measures, which become more involved, particularly for high dimensions; cf. [Sor15].

Some results presented in this paper might be direct consequences of general results in the weak KAM theory; however, we nevertheless provide more explicit information and elementary proofs for the simple example on \mathbb{S}^1 that are particularly connected to the Freidlin–Wentzell theory. For a comprehensive study or survey of the weak KAM theory, we refer to Fathi's book [Fat08] and some very recent books [Sor15, Tra21]. For recent developments of the weak KAM theory in noncompact domains, we refer to [FRF09] for the regularity of Hamiltonians and to [WWY19] for contact Hamiltonian systems where $H(\nabla u(x), u(x), x)$ also depends on u . We particularly refer to [Con01, Gom08, II09, DFIZ16, IS20] for the weak KAM solution as a vanishing discount limit in compact/noncompact domains. The vanishing discount limit of the corresponding optimal control problem does provide a selection principle for weak KAM solutions. See also [CGMT15, MT17], which include a degenerate diffusion term in the vanishing discount limit problem, and see [IMT17] for a duality framework in the vanishing discount problem for fully nonlinear, degenerate elliptic Hamiltonians. The selection principle from the vanishing discount limit is, however, different from the selection principle provided by the large deviation principle for invariant measures. Nonuniqueness of the viscosity solution to the stationary HJE is an important issue even if the Hamiltonian is strictly convex. The nonuniqueness for the vanishing viscosity limit of both stationary HJE and stationary conservation laws are important problems. For instance, constructing a vanishing viscosity approximation to stationary HJE which has a uniform limit is still open [Tra21]. For the stationary transonic flow, multiple stationary entropy shocks were constructed by Steinhoff and Jameson [SJ82], and a selection principle via the stability of the time-dependent problem was studied by Embid, Goodman, and Majda [EGM84] and numerically computed by Shu [Shu88]. We will explain our results below.

For a one-dimensional irreversible diffusion on torus, in Proposition 3.1 we explicitly compute the detailed structures of the Peierls barrier $h(y; x_i)$, which is a central concept in the definition of the projected Aubry set in the Mather–Aubry theory developed in the 1980s. Then we use the Peierls barrier $h(y; x_i)$ to study the detailed structure of the Freidlin–Wentzell variational construction of the rate function

$W^*(x) = W(x) - \min_{x \in \mathbb{S}^1} W(x)$ of the large deviation principle for invariant measures of the diffusion process on \mathbb{S}^1 . This includes two essential steps: (i) the global adjustment for boundary data $W(x_i)$ at the local minimums x_i of the original skew periodic potential $U(x)$, and (ii) the local trimming via a variational representation for $W(x)$; see (1.1) and the local version (3.30).

For step (i), we give an alternative proof in Lemma 3.2 that the variational formula for the boundary data satisfies the discrete weak KAM problem. These boundary data indeed uniquely determine a maximal Lipschitz continuous viscosity solution (see Proposition 4.8) and thus the unique selection principle (3.22) for these boundary data is essential to construct a global energy landscape for the original stochastic process. Particularly, when the original potential $U(x)$ is periodic itself, then we verify that the variational formula for the boundary data must give exactly the same values as the original landscape $U(x)$; see Proposition 3.4. As a byproduct, we also show how to obtain a set of consistent boundary data satisfying the discrete weak KAM problem; see Proposition 4.7.

For step (ii), based on the globally adjusted boundary data and Peierls barriers $h(y; x_i)$, we obtain a local variational representation for $W(x)$, which only depends on the adjacent boundary data and barrier functions; see Proposition 3.3. This local trimming procedure reduces the computations, as shown in the examples in Figure 3. After explaining the variational construction for $W(x)$, in Proposition 3.5, we prove $W(x)$ is a global viscosity solution to

$$(1.4) \quad H(W'(x), x) = W'(x)(W'(x) - U'(x)) = 0, \quad x \in \mathbb{S}^1,$$

satisfying the boundary data uniquely determined via (3.22).

Section 4 focuses on the weak KAM interpretation for $W^*(x) = W(x) - \min W(x)$. We characterize that the projected Aubry set \mathcal{A} is equal to the projected Mather set \mathcal{M}_0 and is equal to all the critical points of $U(x)$. In Corollary 4.5, we prove $W^*(x)$ is a weak KAM solution to (1.4) of negative type, in which the calibrated curves track back to the projected Mather set and those curves are simply solved by the “uphill/downhill” first order ODEs (4.13) and (4.11), respectively. Moreover, the constructed $W(x)$ is the maximal Lipschitz continuous viscosity solution satisfying the boundary data $W(x_i) = W_i$ given in (3.22). These boundary data are chosen via (3.22) and Lemma 3.2, so that $W^*(x) = W(x) - \min_{x \in \mathbb{S}^1} W(x)$ is the rate function for the large deviation principle of the invariant measures of the diffusion process on \mathbb{S}^1 . While all the invariant sets characterized above are the uniqueness sets for the weak KAM solutions to HJE (1.4), there are other uniqueness sets and we show that the uniqueness sets must contain all the local maximums/minimums; see Lemma 4.2. After all these understandings from the weak KAM perspective, we give a probability interpretation for the weak KAM solution $W^*(x)$.

In section 5, we provide more understanding of the obtained weak KAM solution $W^*(x)$, including the exchange of double limits and how one selects a meaningful weak KAM solution that captures the asymptotic behavior of the original stochastic process at each local attractors. In Proposition 5.1, using the property that $W^*(x)$ is an invariant solution to the Lax–Oleinik semigroup representation for the corresponding dynamic HJE, we prove that for a special initial distribution, the large time limit and the zero noise limit can be exchanged for the distribution $\rho_\varepsilon(x, t)$ of the diffusion process, i.e., “ $\lim_{\varepsilon \rightarrow 0} (-\varepsilon \log \lim_{t \rightarrow +\infty} \rho_\varepsilon(x, t)) = \lim_{t \rightarrow +\infty} \lim_{\varepsilon \rightarrow 0} -\varepsilon \log \rho_\varepsilon(x, t)$.” In general, the double limits in both sides exist for any initial data. The right-hand-side limit exists [BS00, Theorem 2.1] but is not unique. However, the left-hand-side limit is unique, which provides a selection principle. That is to say we first take the long time limit

$\lim_{t \rightarrow +\infty} \rho_\varepsilon(x, t)$ which is unique due to ergodicity and then take the zero noise limit $\varepsilon \rightarrow 0$ due to the large deviation principle for invariant measures. In section 5.3, we discuss our selection principle for weak KAM solutions, which are in general not unique; see examples in section 5.1. $W^*(x) = \lim_{\varepsilon \rightarrow 0} (-\varepsilon \log \lim_{t \rightarrow +\infty} \rho_\varepsilon(x, t))$ where the variational formula for boundary data $W(x_i)$ serves as a meaningful selection principle because it is proved to be the rate function of the large deviation principle for the invariant measures. Indeed, the associated viscous HJE computed from the WKB reformulation of the invariant measure $\pi_\varepsilon(x) = e^{-\frac{W_\varepsilon(x)}{\varepsilon}}$ of the irreversible diffusion process is

$$(1.5) \quad W'_\varepsilon(W'_\varepsilon - U') = \varepsilon(W_\varepsilon - U)''.$$

As the rate function of the associated irreversible diffusion process on \mathbb{S}^1 , $W^*(x)$ is unique and can be regarded as the limit of $W_\varepsilon(x)$, in the sense of the large deviation principle (see (2.20)). (1.5) also provides a special construction of a viscosity approximation, which has a uniform vanishing viscosity limit. We point out that in general, the vanishing viscosity approximation method for stationary HJEs only has converged subsequences whose limit is not unique. Our selection principle is different from the widely studied selection principle via the discount limit of the associated optimal control problem in an infinite time horizon. The discount limit method usually cannot capture the long time behavior of the original dynamics. Based on the selection principle in the large deviation sense, the periodic Lipschitz continuous global energy landscape $W^*(x)$ determines the most probable states/path for the original stochastic dynamics as the noise goes to zero; see the generalized Boltzmann analysis through the calibrated curves in section 5.4.

The remaining parts of this paper are organized as follows. In section 2, we introduce the Langevin dynamics on the circle \mathbb{S}^1 and describe the large deviation principle for the invariant measures with both an illustrative example and the abstract result by the Freidlin–Wentzell theory. In section 3, we give explicit properties for the Peierls barriers and use them to prove the variational formula for the global adjustment of boundary data and to construct the global energy landscape $W^*(x)$. The local variational representation, the consistency check, and the viscosity solution property for $W^*(x)$ are given in sections 3.2.3, 3.3, and 3.4, respectively. Section 4 focuses on the characterization of Aubry–Mather sets and the weak KAM solution properties. $W^*(x)$ being the weak KAM solution is proved in section 4.2, and its uniqueness depending on the self-consistent boundary data is discussed in section 4.3. The nonuniqueness of weak KAM solutions and our selection principle, compared with other selection methods, are given in section 5. The probability interpretation for the Freidlin–Wentzell construction of $W^*(x)$ is discussed in section 5.4.

2. The rate function of large deviation principle for the invariant measure of Langevin dynamics on a circle \mathbb{S}^1 . We first introduce a very simple stochastic model, which, however, contains all the representative properties to study the relations between the large deviation principle for invariant measures and the weak KAM theory. This is a one-dimensional irreversible diffusion process on the periodic domain \mathbb{S}^1 , in which the WKB reformulation for the invariant measure $\pi_\varepsilon(x)$ gives a stationary HJE. In section 2.2, we first use a single-well nonperiodic potential $U(x)$ to illustrate the local trimming of the potential $U(x)$ brought by the large deviation principle, and then we describe the general large deviation principle for the invariant measure $\pi_\varepsilon(x)$, which was proved by Freidlin and Wentzell [VF69, VF70]. The associated variational formula for the rate function will be introduced in detail in section 3.

2.1. Langevin dynamics on a circle \mathbb{S}^1 . In this subsection, we first introduce a Langevin dynamics on the simplest closed manifold \mathbb{S}^1 . We start from a Langevin dynamics on a circle \mathbb{S}^1 with a drift in gradient form, i.e., there exists a smooth skew periodic potential $U(x)$ such that $b(x) = -U'(x), x \in \mathbb{S}^1$. This Langevin dynamics on \mathbb{S}^1 reads

$$(2.1) \quad dx = -U'(x)dt + \sqrt{2\varepsilon}dB.$$

Here the skew periodicity of the smooth function $U(x)$ implies there exists a smooth periodic function $\tilde{U}(x)$ such that $U(x) = \tilde{U}(x) - \bar{b}x$ for a constant \bar{b} . Therefore, $U'(x), x \in \mathbb{S}^1$ is a smooth periodic function and

$$(2.2) \quad \int_0^1 U'(x)dx = U(1) - U(0) = -\bar{b}.$$

We refer to the dashed black line in Figure 5 for an example of a skew periodic potential $U(x)$ with three local minimums in one skew period.

The Kolmogorov forward equation corresponding to (2.1) is

$$(2.3) \quad \partial_t \rho_\varepsilon = (\rho_\varepsilon U')' + \varepsilon \rho_\varepsilon'' \quad \text{in } \mathbb{S}^1.$$

Plugging the WKB reformulation $\rho_\varepsilon(x, t) = e^{-\frac{\psi_\varepsilon(x, t)}{\varepsilon}}$ into (2.3) and then taking $\varepsilon \rightarrow 0$, we obtain the dynamic HJE

$$(2.4) \quad \partial_t \psi(x, t) + H(\partial_x \psi(x, t), x) = 0, \quad x \in \mathbb{S}^1,$$

where the Hamiltonian $H : \mathbb{R} \times \mathbb{S}^1 \rightarrow \mathbb{R}$ is

$$(2.5) \quad H(p, x) = p(p - U').$$

Then the corresponding Lagrangian, as the convex conjugate of $H(p, x)$, is given by

$$(2.6) \quad L(s, x) = \sup_{p \in \mathbb{R}} (sp - H(p, x)) = sp^* - H(p^*, x) = \frac{1}{4}(s + U'(x))^2,$$

where p^* solves $s = \partial_p H(p^*, x) = 2p^* - U'(x)$. It is easy to see Hamiltonian $H(p, x)$ is strictly convex w.r.t. p , periodic w.r.t. x while Lagrangian $L(s, x)$ is strictly convex w.r.t. s , periodic w.r.t. x . Another important property is

$$(2.7) \quad L(s, x) \geq 0 \text{ and } L = 0 \text{ if and only if } s = -U'(x).$$

The above ODE flow $\dot{x} = -U'(x)$ can be naturally embedded into the Euler–Lagrangian flow $(x, \dot{x})(t)$ on the tangent bundle $T\mathbb{S}^1$. This special Lagrangian graph $(x, -U'(x))$ enables explicit computations for invariant measures and action minimizing measures/curves; see Mañé [Man92]. So this Lagrangian (2.6) is also known as the Mañé Lagrangian [FR12].

2.2. The invariant measure $\pi_\varepsilon(x)$ and the large deviation principle as $\varepsilon \rightarrow 0$. The corresponding invariant measure $\pi_\varepsilon(x)$ satisfies the stationary Fokker–Planck equation

$$(2.8) \quad \varepsilon \pi_\varepsilon'' + (U' \pi_\varepsilon)' = 0 \quad \text{in } \mathbb{S}^1.$$

Without loss of generality, we assume $\min U(x) = 0$. The unique periodic positive solution π_ε is given by

$$(2.9) \quad \pi_\varepsilon(x) = C_\varepsilon e^{-\frac{U(x)}{\varepsilon}} \int_x^{x+1} e^{\frac{U(y)}{\varepsilon}} dy, \quad x \in \mathbb{S}^1,$$

where C_ε is a normalization constant such that $\int \pi_\varepsilon = 1$. The integral function in (2.9) can be regarded as a corrector to make $\pi_\varepsilon(x)$ periodic. Indeed, recast (2.9) as $\pi_\varepsilon(x) \propto \int_x^{x+1} e^{\frac{U(y)-U(x)}{\varepsilon}} dy$, which is periodic.

If $\bar{b} = 0$, then $U(x)$ is periodic and the above integral in (2.9) is a constant. Thus the Langevin dynamics (2.1) is a reversible process and the periodic invariant measure is directly given by $\pi_\varepsilon(x) \propto e^{-\frac{U(x)}{\varepsilon}}$. Indeed, from (2.9), one can compute the steady flux

$$(2.10) \quad J_\varepsilon = \varepsilon \pi'_\varepsilon + U' \pi_\varepsilon = \varepsilon C_\varepsilon \left(e^{-\frac{\bar{b}}{\varepsilon}} - 1 \right).$$

$\bar{b} = 0$ is equivalent to $J_\varepsilon = 0$ pointwise and thus equivalent to reversibility of the Langevin dynamics (2.1). Then it is obvious that $U(x) = -\varepsilon \log \pi_\varepsilon(x)$ is the rate function in the large deviation principle for the reversible invariant measure $\pi_\varepsilon(x)$.

However, if $\bar{b} \neq 0$, then the Langevin dynamics (2.1) is irreversible and the invariant measure does not have a straightforward formula to serve as a rate function in the large deviation principle. In this case, we define a WKB reformulation

$$(2.11) \quad W_\varepsilon(x) := -\varepsilon \log \pi_\varepsilon(x) = U(x) - \varepsilon \log \int_x^{x+1} e^{\frac{U(y)}{\varepsilon}} dy - \varepsilon \log C_\varepsilon, \quad x \in \mathbb{S}^1.$$

From (2.9), since the solution $\pi_\varepsilon(x)$ to (2.9) has a unique closed formula, $W_\varepsilon(x) = -\varepsilon \log \pi_\varepsilon(x)$ can be uniquely computed up to a constant.

If as $\varepsilon \rightarrow 0$, the limit $W_\varepsilon(x) \rightarrow W^*(x)$ exists for some periodic function $W^*(x)$, then this limit $W^*(x)$ is the rate function for the large deviation principle of the invariant measure $\pi_\varepsilon(x)$. For a peculiar case that $U(x)$ is strictly monotone, then $\pi_\varepsilon(x) = \frac{C}{|U'(x)|} + O(\varepsilon)$ does not have an exponential asymptotic behavior. In this case, $W^*(x) \equiv 0$. Hence we only consider the case when $U(x)$ has minimums.

2.2.1. Illustration of the Laplace principle for a single-well potential $U(x)$. In this subsection, we use the following simple example with a single-well nonperiodic potential $U(x)$ to explicitly compute and simulate the convergence from $W_\varepsilon(x)$ to the globally defined, periodic, Lipschitz continuous rate function $W^*(x)$; see plots of $U(x)$, $W_\varepsilon(x)$, and $W(x)$ in Figure 1.

Take

$$(2.12) \quad U(x) = \cos(2\pi x) - \cos(\pi x) + \frac{9}{8}, \quad x \in [0, 1].$$

Then $U(x)$, $x \in (0, 1)$ is a single basin of attractor of the stable state $x_0 = \frac{1}{\pi} \arccos \frac{1}{4}$, $U_{\min} = U(x_0) = 0$ and the boundary difference is $-\bar{b} = U(1) - U(0) = 2$. One can do skew periodic extension to a C^1 function on \mathbb{R} by $U(x+k) = U(x) - \bar{b}k$, $k \in \mathbb{Z}$. Defining $x^* = \frac{2}{3}$, which has the same value as $U_{\text{exit}} := U(0) = U(x^*)$ for the exit problem, we have

$$0 = U_{\min} < U_{\text{exit}} = U(0) = U(x^*) < U(1) = U_{\max}.$$




FIG. 1. The plot of $U(x)$ in (2.12), $W_\varepsilon(x)$ in (2.15) with different values of ε , and the rate function $W^*(x)$ in (2.17). As $\varepsilon \rightarrow 0$, a uniform convergence from $W_\varepsilon(x)$ to the rate function $W^*(x)$ is shown with a zoom-in plot near the nondifferentiable connection point, where $U(x)$ is cut off from above by the constant U_{exit} . $U(x)$ is skew periodic while both $W_\varepsilon(x)$ and $W(x)$ are periodic.

Since $U_{\max} = U(1)$, $b + U_{\max} = U(0)$ and (2.9) can be reformulated with a different C_ε ,

$$\begin{aligned}
 \pi_\varepsilon(x) &= C_\varepsilon e^{-\frac{U(x)}{\varepsilon}} \left(\int_x^1 e^{\frac{U(z)-U_{\max}}{\varepsilon}} dz + \int_0^x e^{\frac{U(z)-b-U_{\max}}{\varepsilon}} dz \right) \\
 (2.13) \quad &= C_\varepsilon e^{-\frac{U(x)}{\varepsilon}} \left(\int_0^x e^{\frac{U(z)-U(0)}{\varepsilon}} dz + \int_x^1 e^{\frac{U(z)-U(1)}{\varepsilon}} dz \right), \quad x \in \mathbb{S}^1.
 \end{aligned}$$

Since $\min U(x) = 0$, $e^{-\frac{U(x)}{\varepsilon}} = O(1)$ and by the Laplace principle $\int_0^x e^{\frac{U(z)-U(0)}{\varepsilon}} dz + \int_x^1 e^{\frac{U(z)-U(1)}{\varepsilon}} dz \geq O(1)$ as $\varepsilon \rightarrow 0$. Thus $C_\varepsilon \leq O(1)$. When $x \in [0, x^*]$, $U(x) = \min\{U(x), U_{\text{exit}}\}$, then one can directly apply the Laplace principle for the integrals in (2.13). But for $x \in [x^*, 1]$, $U(0) = \min\{U(x), U_{\text{exit}}\}$, so the first integration in (2.13) shall be recast as

$$(2.14) \quad e^{-\frac{U(x)}{\varepsilon}} \int_0^x e^{\frac{U(z)-U(0)}{\varepsilon}} dz = e^{-\frac{U(0)}{\varepsilon}} \int_0^x e^{\frac{U(z)-U(x)}{\varepsilon}} dz.$$

Then from WKB reformulation (2.11), we obtain

$$\begin{aligned}
 (2.15) \quad W_\varepsilon(x) &= -\varepsilon \log \pi_\varepsilon(x) = \min\{U(x), U_{\text{exit}}\} \\
 -\varepsilon \log \left\{ \begin{array}{ll} C_\varepsilon \left(\int_0^x e^{\frac{U(z)-U(0)}{\varepsilon}} dz + \int_x^1 e^{\frac{U(z)-U(1)}{\varepsilon}} dz \right), & x \in [0, x^*]; \\ C_\varepsilon \left(\int_0^x e^{\frac{U(z)-U(x)}{\varepsilon}} dz + e^{\frac{U(0)-U(x)}{\varepsilon}} \int_x^1 e^{\frac{U(z)-U(1)}{\varepsilon}} dz \right), & x \in [x^*, 1]. \end{array} \right.
 \end{aligned}$$

Now every integral term in the above desingularization formula is $O(1)$ and can be numerically implemented. Then by the Laplace principle, we show the last term in (2.15) is an $o(1)$ term,

$$(2.16) \quad \varepsilon \log \begin{cases} C_\varepsilon \left(\frac{1}{2} \sqrt{\frac{2\pi\varepsilon}{|U''(0)|}} + \frac{1}{2} \sqrt{\frac{2\pi\varepsilon}{|U''(1)|}} \right), & x \in [0, x^*]; \\ C_\varepsilon \left(\frac{\varepsilon}{U'(x)} + e^{\frac{U(0)-U(x)}{\varepsilon}} \frac{1}{2} \sqrt{\frac{2\pi\varepsilon}{|U''(1)|}} \right), & x \in [x^*, 1]. \end{cases}$$

Hence, we obtain the rate function $W^*(x)$ for the large deviation principle

$$(2.17) \quad W_\varepsilon(x) \rightarrow W^*(x) := \min\{U(x), U_{\text{exit}}\} \quad \text{in } \mathbb{S}^1.$$

In Figure 1, the WKB reformulation $W_\varepsilon(x)$ is plotted with $\varepsilon = 0.05, 0.01, 0.005, 0.003, 0.002, 0.001$. A uniform convergence from $W_\varepsilon(x)$ to the rate function $W^*(x)$ is shown as $\varepsilon \rightarrow 0$.

For other cases where $U(x)$ has multiple wells, this simple cutoff (local trimming) by a constant as described in the above example is not enough. A globally defined adjustment for the values at each local minimum of $U(x)$ needs to be constructed first, and then the local trimming procedures are applied; see section 3.2. Finding the correct global energy landscape with multiple wells, after proper global adjustment and gluing, is important for the rare events computations; cf. [ELVE19], [GQ12], [ZL16], and [GLLL23]. The global energy landscape correctly measures the action/energy required for the transition from one state to another state, which in general is not the original potential function $U(x)$, as we have already seen from the above simple example (Figure 1).

For the general case, the explicit variational formula for the limit $\lim_{\varepsilon \rightarrow 0} W_\varepsilon(x) = W^*(x) = W(x) - \min W(x)$ was discovered by [VF69, section 8] (see below (3.28)) and we will describe it in detail in section 3. In [VF69, section 8] (see also [FW12, Chapter 6, Theorem 4.3]), this limit $W^*(x)$ is proved to be the rate function for the large deviation principle of $\pi_\varepsilon(x)$ in the following sense. For any $\gamma > 0$, there exists δ_0 such that for any $\delta < \delta_0$, there exists ε_0 such that for any $\varepsilon < \varepsilon_0$

$$(2.18) \quad W(x) - \min W(x) - \gamma \leq -\varepsilon \log \pi_\varepsilon(B_\delta(x)) \leq W(x) - \min W(x) + \gamma.$$

This statement is equivalent to Varadhan's definition [Var16, Definition 2.2] for the large deviation principle on compact domain. Indeed, taking \liminf and \limsup w.r.t. δ and ε , we have

$$(2.19) \quad \begin{aligned} W(x) - \min W(x) - \gamma &\leq \liminf_{\delta \rightarrow 0} \liminf_{\varepsilon \rightarrow 0} -\varepsilon \log \pi_\varepsilon(B_\delta(x)) \\ &\leq \limsup_{\delta \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} -\varepsilon \log \pi_\varepsilon(B_\delta(x)) \leq W(x) - \min W(x) + \gamma. \end{aligned}$$

Thus since γ is arbitrary,

$$(2.20) \quad W(x) - \min W(x) = \liminf_{\delta \rightarrow 0} \liminf_{\varepsilon \rightarrow 0} -\varepsilon \log \pi_\varepsilon(B_\delta(x)) = \limsup_{\delta \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} -\varepsilon \log \pi_\varepsilon(B_\delta(x)).$$

Thus, (2.19) is exactly the lower bound and upper bound estimates in [Var16, Definition 2.2] and thus implies the large deviation principle for the invariant measure $\pi_\varepsilon(x)$ with the rate function $W(x) - \min W(x)$. In one dimension, [FG12] provides a direct

proof for the limit of $-\varepsilon \log \pi_\varepsilon(x)$ via Laplace's principle and recovers the variational formula (3.28).

We remark that for general stochastic processes, for instance, for the large deviation principle in the thermodynamic limit for the invariant measures of the chemical reaction process [GL23], one can also directly study the upper semicontinuous (USC) envelope of the WKB reformulation $W_\varepsilon(x)$

$$(2.21) \quad \overline{W}(x) := \limsup_{\varepsilon \rightarrow 0, x_j \rightarrow x} W_\varepsilon(x_j)$$

and the lower semicontinuous (LSC) envelope of $W_\varepsilon(x)$

$$(2.22) \quad \underline{W}(x) := \liminf_{\varepsilon \rightarrow 0, x_j \rightarrow x} W_\varepsilon(x_j).$$

By these definitions, if the large deviation principle (2.20) holds, then necessarily one obtains

$$(2.23) \quad \underline{W}(x) \leq W(x) - \min W(x) \leq \overline{W}(x).$$

In [GL23, Proposition 4.1], under the detailed balance condition for the chemical reaction process, the USC envelope $\overline{W}(x)$ is proved to be a USC viscosity solution to the corresponding stationary HJE in the Barron–Jensen sense [BJ90].

2.2.2. Variational formula for $W^*(x)$ through Varadhan's lemma. Recall the WKB reformulation $W_\varepsilon(x)$ of invariant measure $\pi_\varepsilon(x)$. From the large deviation principle (2.20) for the invariant measure $\pi_\varepsilon(x)$, Varadhan's lemma [Var66, Var16] provides another variational formula for the rate function $W^*(x)$. Below, we carry out details for this formula.

Using Varadhan's lemma [Var16, Theorem 2.5], we know for any test function $f \in C(\mathbb{S}^1)$,

$$(2.24) \quad \sup_{y \in \mathbb{S}^1} (f(y) - W^*(y)) = \lim_{\varepsilon \rightarrow 0} -\varepsilon \log \int_{\mathbb{S}^1} e^{\frac{f(y)}{\varepsilon}} \pi_\varepsilon(y) dy.$$

Denote the integral above as A_ε and compute it via the closed formula (2.11)

$$(2.25) \quad \begin{aligned} A_\varepsilon &:= \int_{\mathbb{S}^1} e^{\frac{f(y)}{\varepsilon}} \pi_\varepsilon(y) dy = C_\varepsilon \int_0^1 e^{\frac{f(y)-U(y)}{\varepsilon}} \int_y^{y+1} e^{\frac{U(z)}{\varepsilon}} dz dy \\ &= C_\varepsilon \int_0^1 \int_0^z e^{\frac{f(y)-U(y)+U(z)}{\varepsilon}} dy dz + C_\varepsilon e^{-\frac{b}{\varepsilon}} \int_0^1 \int_z^1 e^{\frac{f(y)-U(y)+U(z)}{\varepsilon}} dy dz. \end{aligned}$$

Here we used exchange order of integrals and skew periodicity (2.2). However, there is no simple way to directly study this globally defined limiting function. Thus we go back to the Freidlin–Wentzell variational construction below.

In the next section, under the assumption that there are finite critical points x_i for $U(x)$, we reinterpret and give an alternative proof for the construction of a global periodic energy landscape $W(x)$ from locally defined quasi-potential after adjusting levels and then proper trimming and gluing. From the formula of $W_\varepsilon(x)$ in (2.11), the limiting rate function $W^*(x)$ can be viewed as the original potential $U(x)$ with an additional corrector computed from the Laplace principle for $\varepsilon \log \int_x^{x+1} e^{\frac{U(y)}{\varepsilon}} dy$. This provides the recipes to (i) globally adjust the levels $W(x_i)$ of each quasi-potential for each stable basin via (3.22) and then (ii) to locally trimming from above and

glue to construct $W^*(x) = W(x) - \min_i W(x_i)$ via (3.28) or (3.30). This global energy landscape is continuous and is proved to be the rate function of the large deviation principle for the invariant measure of Langevin dynamics on a circle \mathbb{S}^1 [FW12, Chapter 6, Theorem 4.3]. We will also prove that $W^*(x)$ is a viscosity solution to HJE (3.48); see Proposition 3.5.

3. Freidlin–Wentzell variational construction of periodic Lipschitz continuous global energy landscape $W^*(x)$. In this section, we focus on the detailed description of the Freidlin–Wentzell variational construction of the rate function $W^*(x)$; see (3.24) and (3.28). Using the one-dimensional irreversible diffusion on torus (2.3), we give an alternative elementary proof for the variational formula (3.24) for determining the boundary data and elaborating explicit properties of $W^*(x)$ such as the shape of the nondifferentiable part. Those boundary values are globally defined and are the most crucial ingredient to obtain the unique, Lipschitz continuous, periodic global energy landscape that can correctly represent the exponentially small probability in the large deviation principle. Before giving the variational formula, we revisit and prove the detailed characterizations of two essential concepts of least action functions: the quasi-potential (a.k.a. the Mañé potential) and the Peierls barrier. Based on these explicit properties, we then give the construction of a global energy landscape in section 3.2 based on (i) the global adjustment for boundary data on the local minimums and (ii) a local trimming procedure via adjacent boundary data and the Peierls barrier; see Lemma 3.2 and Proposition 3.3. Last, we give a consistent verification to show the variational representation of $W^*(x)$ is indeed reduced to the original potential function $U(x)$ if the diffusion process is reversible; see Proposition 3.4.

Let us first clarify that we always work on the case that $U(x)$ has finite many critical points indexed as follows. Assume there are k stable local minimums

$$(3.1) \quad x_1, x_2, \dots, x_k$$

of $U(x)$, interleaved by k unstable local maximums

$$(3.2) \quad x_{\frac{1}{2}}, x_{1+\frac{1}{2}}, \dots, x_{k+\frac{1}{2}} = x_{\frac{1}{2}} + 1.$$

Without loss of generality, we assume

$$(3.3) \quad 0 = x_{\frac{1}{2}} < x_1 < x_{1+\frac{1}{2}} < x_2 < \dots < x_k < x_{k+\frac{1}{2}} = 1.$$

Denote other duplicated points outside $[0, 1]$ as $x_{i+\ell k} = x_i + \ell \in \mathbb{R}$ for any $\ell \in \mathbb{Z}$.

3.1. Properties of Peierls barrier $h(y; x_i)$ and Mañé potential $v(y; x_i)$.

In this subsection, we revisit two essential concepts of least action functions: the quasi-potential (a.k.a. the Mañé potential) and the Peierls barrier. In our one-dimensional example, we further explore the explicit shape characterizations for those least action functions, which are important properties for the later construction of global energy landscape.

3.1.1. Quasi-potentials and critical Mañé value $c^* = 0$. Quasi-potential is an essential concept introduced in the Freidlin–Wentzell theory, while the local quasi-potential within a basin of attraction of stable states is widely used for computing the barrier of exit problems for stochastic processes. We explain below the globally defined quasi-potential, which is now called the Mañé potential following the convention in the weak KAM theory.

Given any starting point x_0 , not necessarily critical points, the Mañé potential is defined as

$$(3.4) \quad v(y; x_0) := \inf_{T \geq 0, \gamma(0)=x_0, \gamma(T)=y} \int_0^T (L(\dot{\gamma}(t), \gamma(t)) + c) dt.$$

It is well known that the critical Mañé value for the Lagrangian (2.6) is zero $c^* = 0$. So from now on, we drop c in the definition of the Mañé potential. As it is an essential concept, we provide descriptions of four characterizations for c^* below.

(i) The definition of the critical Mañé value is (cf. [CI99])

$$(3.5) \quad c^* = \sup\{c \in \mathbb{R}; \exists \text{ closed curve } x(\cdot) \text{ s.t. } \int_0^T (L(\dot{x}(t), x(t)) + c) dt < 0\}.$$

Since $L \geq 0$, we know at least $c^* \leq 0$. On the other hand, if $c^* < 0$, then one can choose a standing curve $x(t) \equiv x_i$ at a steady state x_i such that $\dot{x} = -U'(x) \equiv 0$. Then one has $L \equiv 0$ while $\int_0^T (L(\dot{x}(t), x(t)) + c^*) dt < 0$. Thus $c^* = 0$.

(ii) From [Fat08, Definition 4.2.6], one can verify

$$(3.6) \quad c^* := \inf\{c \in \mathbb{R}; \text{ there exists } u \in \text{Lip}(\mathbb{S}^1) \text{ s.t. } H(u'(x), x) \leq c \text{ a.e.}\} = 0.$$

Indeed, on the one hand, taking $u \equiv 0$ implies $H(u'(x), x) = 0$, so at least $c^* \leq 0$. On the other hand, since $H = (u' - \frac{U'}{2})^2 - (\frac{U'}{2})^2$, for any $u \in \text{Lip}(\mathbb{S}^1)$, $H \geq 0$ at critical point x_i of U . Thus c^* cannot be negative, so $c^* = 0$.

There are two other methods for characterizing c^* : (iii) c^* can also be computed via the min-max problem (cf. [Eva08, Theorem 4.1]),

$$c^* = - \inf_{\varphi \in C^1(\mathbb{S}^1)} \max_{x \in \mathbb{S}^1} \varphi'(x) (\varphi'(x) - U'(x)) = 0;$$

(iv) c^* can be computed via action minimizing (Mather) measures (cf. [Eva08, Theorem 2.7]): Let $\mathcal{P}_{\text{inv}}(\mathbb{R} \times \mathbb{S}^1)$ be the collection of probability measures μ on $\mathbb{R} \times \mathbb{S}^1$ that is invariant under the Lagrangian flow. Then

$$c^* = - \inf_{\mu \in \mathcal{P}_{\text{inv}}} \int_{\mathbb{R} \times \mathbb{S}^1} L(s, x) d\mu(s, x) = 0.$$

See also [Man96] for a relaxed minimization which relaxes the invariant Lagrangian flow condition. The measure achieving the minimum is called the Mather measure. There are many Mather measures μ on $\mathbb{R} \times \mathbb{S}^1$ for our problem. For instance, we take $\mu = \delta(s)\delta(x - x_i)$, where x_i is a steady state, and it is easy to verify the minimum $c^* = 0$ is achieved.

3.1.2. Characterization of Peierls barriers $h(y; x_i)$ on \mathbb{S}^1 . We point out that for the above case that the starting point x_i is a stable/unstable critical point of $U(x)$, another important concept, called the Peierls barrier, is defined as

$$(3.7) \quad h(y; x_i) := \liminf_{T \rightarrow +\infty} \inf_{\gamma(0)=x_i, \gamma(T)=y} \int_0^T L(\dot{\gamma}(t), \gamma(t)) dt.$$

From the computations for left/right Mañé values in (3.10) and (3.12), it is easy to see for x_i being a critical point of $U(x)$, then $v(y; x_i) = h(y; x_i)$ for any $y \in \mathbb{S}^1$. Thus from now on, we use Peierls barrier $h(y; x_i)$ instead of $v(y; x_i)$ whenever the starting point is a critical point of $U(x)$.

Before we explain explicitly the global energy landscape $W^*(x)$ construction, we characterize the explicit formula for Peierls barriers $h(y; x_i)$ with the specific nondifferential point, connection shape, and periodicity; see Figure 5. This will also serve as a key observation for justifying the weak KAM solution later. In this paper, we always assume the orientation of $x \in \mathbb{S}^1$ belonging to an interval $x \in (a, b)$ is counterclockwise.

PROPOSITION 3.1. *Assume there are k stable local minimums of $U(x)$, interleaved by k unstable local maximums indexed as (3.1) and (3.2). Then*

- (i) *the Peierls barrier $h(y; x_i) \geq 0$ is Lipschitz continuous and periodic;*
- (ii) *there exists $x^* \in \mathbb{S}^1$ such that $h(y; x_i)$ is nonincreasing in (x^*, x_i) to zero and then nondecreasing in $(x_i, x^* + 1)$ back to the same level $h(x^*; x_i) = h(x^* + 1; x_i)$;*
- (iii) *the only one possible nondifferential point is the connection point x^* , where either an increasing function is connected to a constant or a constant is connected to a decreasing function, that is to say, $h(y; x_i)$ is a C^1 function cut off at most once by a constant from above.*
- (iv) *$h(y; x_i)$ is the maximal Lipschitz continuous viscosity solution to HJE*

$$(3.8) \quad H(h'(y), y) = h'(h' - U') = 0, \quad y \in \mathbb{S}^1,$$

and satisfies $h(x_i; x_i) = 0$.

This proposition on the characterization of Peierls barriers is basically known in the weak KAM theory but here we give detailed properties on the periodicity and explicit shape of $h(y; x_i)$.

Proof. First, we define a right barrier function for $y \in [x_i, x_{i+k}] \subset \mathbb{R}$

$$(3.9) \quad h_R(y; x_i) := \inf_{T \geq 0, \gamma(0) = x_i, \gamma(T) = y} \int_0^T \frac{1}{4} |\dot{\gamma} + U'(\gamma)|^2 dt$$

for an exit problem starting from x_i to the right until the point x_{i+k} passing through several local minimums $x_{i+1}, \dots, x_{i+k-1} \in \mathbb{R}$. To see explicitly the formula for the barrier function, from each stable minimum x_i to its adjacent critical points, we can first compute

$$(3.10) \quad \begin{aligned} h_R(x_{i+\frac{1}{2}}; x_i) &= \inf_{T \geq 0, \gamma(0) = x_i, \gamma(T) = x_{i+\frac{1}{2}}} \int_0^T \frac{1}{4} |\dot{\gamma} + U'(\gamma)|^2 dt \\ &= \inf_{T \geq 0, \gamma(0) = x_i, \gamma(T) = x_{i+\frac{1}{2}}} \int_0^T \left(\frac{1}{4} |\dot{\gamma} - U'(\gamma)|^2 + \dot{\gamma} U' \right) dt \geq U(x_{i+\frac{1}{2}}) - U(x_i). \end{aligned}$$

Here the equality holds if and only if $\dot{\gamma} = U'(\gamma)$ and $\gamma(+\infty) = x_{i+\frac{1}{2}}$, so $h_R(x_{i+\frac{1}{2}}; x_i) = U(x_{i+\frac{1}{2}}) - U(x_i)$. It is usually referred as the “uphill” path from x_i to $x_{i+\frac{1}{2}}$; cf. [FW12]. Similarly, the left barrier from x_i to the left to $x_{i-\frac{1}{2}}$ is

$$(3.11) \quad h_L(x_{i-\frac{1}{2}}; x_i) := \inf_{T \geq 0, \gamma(0) = x_i, \gamma(T) = x_{i-\frac{1}{2}}} \int_0^T \frac{1}{4} |\dot{\gamma} + U'(\gamma)|^2 dt = U(x_{i-\frac{1}{2}}) - U(x_i).$$

Conversely, for the “downhill” path starting from $x_{i+\frac{1}{2}}$ along $\dot{\gamma} = -U'(\gamma)$ and $\gamma(+\infty) = x_{i+1}$, we have

$$(3.12) \quad h_R(x_{i+1}; x_{i+\frac{1}{2}}) = \inf_{T \geq 0, \gamma(0)=x_{i+\frac{1}{2}}, \gamma(T)=x_{i+1}} \int_0^T \frac{1}{4} |\dot{\gamma} + U'(\gamma)|^2 dt = 0.$$

Thus

$$(3.13) \quad \begin{aligned} h_R(x_{i+1}; x_i) &= h_R(x_{i+\frac{1}{2}}; x_i) = U(x_{i+\frac{1}{2}}) - U(x_i); \\ h_L(x_{i-1}; x_i) &= h_L(x_{i-\frac{1}{2}}; x_i) = U(x_{i-\frac{1}{2}}) - U(x_i). \end{aligned}$$

Other barriers passing through multiple wells can be computed repeatedly.

Thus the barrier formula h_R for this one-dimensional least action problem (multiple exit problems) is given by a least action problem for the piecewise C^1 curve connecting x_i to $y \in [x_i, x_{i+k}]$,

$$(3.14) \quad h_R(y; x_i) = \begin{cases} U(y) - U(x_i), & y \in [x_i, x_{i+\frac{1}{2}}], & \text{increase;} \\ U(x_{i+\frac{1}{2}}) - U(x_i), & y \in [x_{i+\frac{1}{2}}, x_{i+1}], & \text{constant;} \\ U(x_{i+\frac{1}{2}}) - U(x_i) + U(y) - U(x_{i+1}), & y \in [x_{i+1}, x_{i+\frac{3}{2}}], & \text{increase;} \\ \dots & \dots & \dots \\ \sum_{j=i}^{i+k-1} [U(x_{j+\frac{1}{2}}) - U(x_j)], & y \in [x_{i+k-\frac{1}{2}}, x_{i+k}], & \text{constant.} \end{cases}$$

We emphasize $U(x)$ is skew periodic function defined on the whole \mathbb{R} , so $h_R(y; x_i)$ is well-defined. It's easy to see $h_R(y; x_i)$ is a nondecreasing $C^1([x_i, x_{i+k}])$ function.

Similarly, a nonincreasing $C^1([x_{i-k}, x_i])$ function $h_L(y; x_i), y \in [x_{i-k}, x_i]$ can be computed to serve as a left barrier function for the exit problem starting from x_i to the left until the point x_{i-k} ,

$$(3.15) \quad h_L(y; x_i) = \begin{cases} \sum_{j=i-k+1}^i [U(x_{j-\frac{1}{2}}) - U(x_j)], & y \in [x_{i-k}, x_{i-k+\frac{1}{2}}], & \text{constant;} \\ \dots & \dots & \dots \\ U(x_{i-\frac{1}{2}}) - U(x_i) + U(y) - U(x_{i-1}), & y \in [x_{i-\frac{3}{2}}, x_{i-1}], & \text{decrease;} \\ U(x_{i-\frac{1}{2}}) - U(x_i), & y \in [x_{i-1}, x_{i-\frac{1}{2}}], & \text{constant;} \\ U(y) - U(x_i), & y \in [x_{i-\frac{1}{2}}, x_i], & \text{decrease.} \end{cases}$$

For other points, due to screw periodicity of $U(x)$, one can naturally define

$$h_L(y \pm 1, x_{i \pm k}) = h_L(y, x_i), \quad h_R(y \pm 1, x_{i \pm k}) = h_R(y, x_i).$$

See Figure 2 for the illustration of the left barrier $h_L(y; x_i)$ and the right barrier $h_R(y; x_i)$.

Second, since $U'(x)$ is periodic, for $y \in \mathbb{S}^1$, we compute the Peierls barrier

$$(3.16) \quad h(y; x_i) := \inf_{T \geq 0, \gamma(0)=x_i, \gamma(T)=y} \int_0^T \frac{1}{4} |\dot{\gamma} + U'(\gamma)|^2 dt.$$

Using h_R, h_L , we represent $h(y; x_i), y \in [0, 1]$, as

$$(3.17) \quad h(y; x_i) = \begin{cases} \min\{h_L(y; x_i), h_R(y+1; x_i)\}, & y \leq x_i, \\ \min\{h_L(y-1; x_i), h_R(y; x_i)\}, & y > x_i, \end{cases}$$

which will be proved below to be a periodic, Lipschitz continuous function. We first give a key observation, which will be used in the characterization of the shape of




FIG. 2. The construction of the Peierls barrier $h(y; x_i)$ via the left barrier $h_L(y; x_i)$ and right barrier $h_R(y; x_i)$. The dashed black line is the skew periodic potential $U(x)$ with three local minimums 1, 0, 2 in $\mathbb{S}^1 = [6_L, 6]$. The solid green line is the left barrier starting from $x_i = 3$ and is nondecreasing clockwise. The solid purple line is the right barrier starting from $x_i = 3$ and is nondecreasing counterclockwise. After finding x^* such that $h_L(x^*; x_i) = h_R(x^* + 1; x_i)$, $h(y; x_i)$ is periodic, Lipschitz continuous with only one cutoff by a constant from the above at the nondifferential point x^* .

the global energy landscape as well. Notice $[x_i, x_{i+\frac{1}{2}}] = [x_{i-k}, x_{i-k+\frac{1}{2}}] + 1$ is on the increasing interval of the i th well $U(x)$, where h_R is increasing and h_L is a constant. Similarly, $[x_{i-\frac{1}{2}}, x_i] = [x_{i+k-\frac{1}{2}}, x_{i+k}] - 1$ is on the decreasing interval of the i th well $U(x)$, where h_R is a constant and h_L is decreasing. Below, we characterize $h(y; x_i)$.

Step 1. Since $h_R(y; x_i)$ is nondecreasing and $h_L(y; x_i)$ is nonincreasing, there always exists x^* such that $h_L(x^*; x_i) = h_R(x^* + 1; x_i)$. Therefore, for $x^* \leq y \leq x_i$, the minimum (3.17) is attained at $h_L(x)$, while for $x_i \leq y \leq x^* + 1$, the minimum (3.17) is attained at $h_R(x)$. Thus the Peierls barriers is given by

$$(3.18) \quad h(y; x_i) = \begin{cases} h_L(y; x_i), & x^* \leq y \leq x_i, \\ h_R(y; x_i), & x_i \leq y \leq x^* + 1. \end{cases}$$

Immediate consequences are that $h(y; x_i)$ is a C^1 function in $(x^*, x^* + 1)$, nonincreasing in (x^*, x_i) to zero, and nondecreasing in $(x_i, x^* + 1)$ back to the same level

$$h(x^*; x_i) = h(x^* + 1; x_i).$$

Thus $h(y; x_i)$ has continuously periodic extension and the only possible nondifferential point is $x^* + \mathbb{Z}$; see Figure 2 for the construction of $h(y; x_i)$ via $h_L(y; x_i), h_R(y; x_i)$. Hence we obtained conclusions (i) and (ii).

Step 2. We prove the type of the nondifferentiability for point x^* .

First, there exists $i - k \leq \ell \leq i$ such that $x^* \in [x_{\ell-\frac{1}{2}}, x_{\ell+\frac{1}{2}}]$. We only need to consider three cases. Case (1): $x^* = x_\ell$ or $x_{\ell \pm \frac{1}{2}}$, then $h'(x^*; x_i) = h'_L(x^*) = h'_R(x^*) = 0$ is differentiable. Case (2): if $x^* \in (x_\ell, x_{\ell+\frac{1}{2}})$, then from the formula (3.18), we know $h'(x^*_+; x_i) = h'_L(x^*_+; x_i) = 0$ while $h'(x^*_-; x_i) = h'_R(x^*_-; x_i) = U'(x^*) > 0$. This case implies that an increasing function is connected to a constant at the nondifferential point x^* . Case (3): if $x^* \in (x_{\ell-\frac{1}{2}}, x_\ell)$, then the left derivative $h'(x^*_-; x_i) = 0$ while

the right derivative exists and is negative $h'(x_+^*; x_i) = U'(x^*) < 0$. This implies that a constant is connected to a decreasing function at the nondifferential point x^* . Therefore, we obtained the conclusion (iii).

Step 3. One can directly verify $h(y; x_i)$ is a viscosity solution based on the definition in [BD+97, p. 5]. The maximality of $h(y; x_i)$ follows [Tra21, Theorem 2.41] or [FRF09, Theorem 2.4] only with small modifications. Let $\tilde{u}(y)$ be a Lipschitz continuous viscosity subsolution to (3.8) satisfying $\tilde{u}(x_i) = 0$ and thus an almost everywhere subsolution satisfying $H(\nabla\tilde{u}(x), x) \leq 0$ a.e. $x \in \mathbb{S}^1$. So for any absolutely continuous curve $\gamma(\cdot)$ with $\gamma(0) = x_i$ and $\gamma(t) = y \in \mathbb{S}^1$, we have

$$\begin{aligned} \tilde{u}(y) - \tilde{u}(x_i) &= \int_0^t \nabla\tilde{u}(\gamma(s)) \cdot \dot{\gamma}(s) ds \\ &\leq \int_0^t (L(\dot{\gamma}(s), \gamma(s)) + H(\nabla\tilde{u}(\gamma(s)), \gamma(s))) ds \leq \int_0^t L(\dot{\gamma}(s), \gamma(s)) ds. \end{aligned}$$

Then taking infimum w.r.t. γ and \liminf w.r.t. t , we obtain

$$(3.19) \quad \tilde{u}(y) \leq \liminf_{t \rightarrow +\infty} \inf_{\gamma; \gamma(0)=x_i, \gamma(t)=y} \int_0^t L(\dot{\gamma}(s), \gamma(s)) ds = h(y; x_i).$$

Thus the Peierls barrier $h(y; x_i)$ is the maximal Lipschitz continuous viscosity solution satisfying (3.8). \square

Remark 1. The shape of the Peierls barrier $h(y; x_{i+\frac{1}{2}})$ starting from the local maximums $x_{i+\frac{1}{2}}$ can be characterized with the same arguments. The only difference is $h(y; x_{i+\frac{1}{2}}) = 0$ for $y \in [x_i, x_{i+1}]$. Then outside $[x_i, x_{i+1}]$, one can use $h_R(y; x_{i+1})$ and $h_L(y; x_i)$ to construct $h(y; x_{i+\frac{1}{2}})$.

3.2. Freidlin–Wentzell variational construction for the rate function $W(x)$ via boundary values W_i at stable states and Peierls barriers $h(y; x_i)$. Based on the previous explicit characterization of Peierls barriers starting from each stable state, in this subsection, we describe and give an alternative proof for the Freidlin–Wentzell variational formula for determining the boundary values. Those boundary values are globally defined and are the most crucial ingredient to obtain the unique, Lipschitz continuous, periodic global energy landscape that can correctly represent the exponentially small probability in the large deviation principle. After obtaining the global adjustment of boundary values, the variational construction for the rate function $W^*(x)$ is indeed a local trimming procedure; see section 3.2.3 for the local representation of $W(x)$.

3.2.1. Determine boundary values W_j on stable states. Now we determine the boundary values $W(x_i)$ at stable minimum x_i . For any $j = 1, \dots, k$, recall $h_R(x_i; x_{j+1})$ defined in (3.14). To compute a counterclockwise path connecting $x_{j+1} \in \mathbb{S}^1$ to $x_i \in \mathbb{S}^1$, we introduce a tilde notation for the total cost of this path on \mathbb{S}^1 ,

$$(3.20) \quad \tilde{h}_R(x_i; x_{j+1}) := \begin{cases} h_R(x_i; x_{j+1}), & j < i, \\ h_R(x_{i+k}; x_{j+1}), & j \geq i. \end{cases}$$

Similarly, using $h_L(x_i; x_j)$ defined in (3.15), the total cost for a clockwise path connecting $x_j \in \mathbb{S}^1$ to $x_i \in \mathbb{S}^1$ is

$$(3.21) \quad \tilde{h}_L(x_i; x_j) := \begin{cases} h_L(x_i; x_j), & j \geq i, \\ h_L(x_{i-k}; x_j), & j < i. \end{cases}$$




FIG. 3. Two examples for $W^*(x)$ computed from the variational formula (3.28) with different consistent boundary data. (Left) The boundary values W_1, W_2, W_3 satisfy the Freidlin–Wentzell variational formula for boundary data (3.22). The dashed line is the original potential $U(x)$ and one can see the global adjustment for W_1, W_2, W_3 . The explicit formula for $W^*(x)$ in (5.1) is marked with a gray strip. (Right) $W^*(x)$ constructed with $W(x_1) = W(x_2) = W(x_3) = 0$ as boundary data, which does not satisfy (3.22). If one adds an additional boundary condition $W(x_{1+\frac{1}{2}}) = 0$, then one obtains a different solution which vanishes in the interval $[x_1, x_2]$, shown as dashed line. Both figures use the same Peierls barriers: $h(y; x_1)$ shown in a red line, $h(y; x_2)$ shown in a green line, and $h(y; x_3)$ shown in an orange line. At each connection point, only two lifted Peierls barriers turn on to finish the local trimming procedures, as described in Proposition 3.3.

Then following [FW12, Chapter 6, equation (4.2)], define

$$(3.22) \quad W_i := \min_{j=1, \dots, k} (\tilde{h}_R(x_i; x_{j+1}) + \tilde{h}_L(x_i; x_j)), \quad i = 1, \dots, k.$$

We refer to the example in Figure 3 (left) for a globally adjusted boundary data W_i , $i = 1, 2, 3$, satisfying (3.22) and the construction of $W^*(x)$ based on those boundary data. With these specially adjusted boundary data, $W^*(x)$ is proved to be the rate function of the large deviation principle for invariant measures (2.18) [FW12, Chapter 6, Theorem 4.3]. We also provide a coarse-grained Markov chain interpretation in Appendix C.

Notice all x_i are stable critical points, so the explicit formula for the Peierls barrier $h(x_i; x_j)$ in (3.17) is recast as

$$(3.23) \quad h(x_i; x_j) = \min\{\tilde{h}_R(x_i; x_j), \tilde{h}_L(x_i; x_j)\}, \quad i, j = 1, \dots, k.$$

In the following lemma, we prove the boundary data satisfying the variational formula (3.22) is indeed a consistent data set satisfying the discrete weak KAM problem (3.24).

LEMMA 3.2. *Let $h(x_i; x_j)$ be the Peierls barrier. The values of W_i , $i = 1, \dots, k$, defined in (3.22) solves the discrete weak KAM problem*

$$(3.24) \quad W_i = \min_{j=1, \dots, k} \{W_j + h(x_i; x_j)\} \quad \forall i = 1, \dots, k.$$

Proof. To verify (3.24), it is sufficient to verify that for any ℓ , there exists m such that

$$(3.25) \quad W_i \leq \tilde{h}_R(x_i; x_{m+1}) + \tilde{h}_L(x_i; x_m) \leq \tilde{h}_R(x_j; x_{\ell+1}) + \tilde{h}_L(x_j; x_\ell) + h(x_i, x_j).$$

Indeed, taking the minimum in ℓ , we have $W_i \leq W_j + h(x_i, x_j)$, and then taking the minimum in j , we have $W_i \leq \min_{j=1, \dots, k} \{W_j + h(x_i; x_j)\}$. Particularly, the equality holds for $j = i$.

Now we prove (3.25) for the case $h(x_i; x_j) = \min\{\tilde{h}_R(x_i; x_j), \tilde{h}_L(x_i; x_j)\} = \tilde{h}_L(x_i; x_j)$, and the other one has the same argument.

- (i) If $\ell \in \{j, j+1, \dots, i-1\}$ is on the counterclockwise path from x_j to x_i , then taking $m = \ell$, we obtain

$$(3.26) \quad \tilde{h}_R(x_i; x_{\ell+1}) + \tilde{h}_L(x_i; x_\ell) \leq \tilde{h}_R(x_j; x_{\ell+1}) + \tilde{h}_L(x_j; x_\ell) + \tilde{h}_L(x_i; x_j)$$

due to $\tilde{h}_R(x_i; x_{\ell+1}) \leq \tilde{h}_R(x_j; x_{\ell+1})$ and $\tilde{h}_L(x_i; x_\ell) \leq \tilde{h}_L(x_j; x_\ell) + \tilde{h}_L(x_i; x_j)$.

- (ii) If $\ell \in \{i, i+1, \dots, j-1\}$ is on the clockwise path from x_j to x_i , then taking $m = i-1$, since $\tilde{h}_R(x_i; x_i) = 0$ and $\tilde{h}_L(x_i; x_{i-1}) \leq \tilde{h}_L(x_j; x_\ell) + \tilde{h}_L(x_i; x_j)$, we have

$$(3.27) \quad \tilde{h}_R(x_i; x_i) + \tilde{h}_L(x_i; x_{i-1}) \leq \tilde{h}_R(x_j; x_{\ell+1}) + \tilde{h}_L(x_j; x_\ell) + \tilde{h}_L(x_i; x_j).$$

Thus (3.25) is proved, and so is the lemma. \square

3.2.2. Variational construction for $W^*(x)$ via boundary values W_i on stable states x_i . With the above boundary values W_i , $i = 1, \dots, k$, on all the stable minima, the global energy landscape is defined as [FW12, Chapter 6, Theorem 4.3]

$$(3.28) \quad W(x) = \min_{j=1, \dots, k} \{W_j + h(x; x_j)\} \quad \forall x \in \mathbb{S}^1.$$

Later in section 4, we will prove that $W(x)$ is indeed a weak KAM solution to the HJE (3.48). We also characterize the corresponding projected Aubry set \mathcal{A} in section 4.1. After including the induced boundary values on other critical point (local maximums) in \mathcal{A} , this $W(x)$ satisfies the usual representation (cf. [Tra21, Theorem 7.4]) via the boundary data on the projected Aubry set for the weak KAM solution; see Lemma 4.3.

We remark that the boundary values W_i to the discrete weak KAM problem (3.24) are not uniquely determined because $W_i = 0$, $i = 1, \dots, k$, are also admissible boundary values satisfying (3.24); see Figure 3 (right), for instance. Meanwhile, a constant shift of W_i is also a solution to (3.24). We refer to section 5 for examples of nonuniqueness.

However, the construction described above using the uniquely determined boundary data W_i and the trimming of $W_i + h(x; x_i)$ has clear probability meaning via the large deviation principle for the invariant measure $\pi_\varepsilon(x)$. From [FW12, Chapter 6, Theorem 4.3],

$$(3.29) \quad W^*(x) = W(x) - \min_i W(x_i)$$

gives the rate function in the large deviation principle for the invariant measure $\pi_\varepsilon(x)$ to the Langevin dynamics (2.1) on \mathbb{S}^1 . In section 4, we will explore more properties of $W(x)$ from the weak KAM viewpoint and use the corresponding projected Aubry–Mather set to give a probability interpretation of the global energy landscape $W^*(x)$.

3.2.3. Local representation for $W(x)$. Based on the globally adjusted boundary values W_j , the rate function $W(x)$ can be constructed in (3.28). In the following proposition, we show that the variational formula (3.28) indeed has a local representation depending only on the boundary values of the adjacent local minima and barrier functions. This procedure is thus also referred to as a local trimming procedure. We refer to Figure 3 for an illustration of the local trimming.

PROPOSITION 3.3. *Let $W(x)$ be given by (3.28) with boundary values W_j , $j = 1, \dots, k$. Assume the boundary values W_j satisfy the discrete weak KAM problem (3.24). Then $W(x)$ has a local representation that, for any $x \in [x_i, x_{i+1}]$ for some $i = 1, \dots, k$,*

$$(3.30) \quad W(x) = \min\{W_i + h_R(x; x_i), W_{i+1} + h_L(x; x_{i+1})\},$$

where $h_R(x; x_i)$ and $h_L(x; x_{i+1})$ are the locally defined, right/left barrier functions in (3.14) and (3.15). Consequently, at each local maximum $x_{i+\frac{1}{2}}$, there is at most one flat connection either on the left of $x_{i+\frac{1}{2}}$ or on the right of $x_{i+\frac{1}{2}}$.

Proof. Assume $W(x)$ defined in (3.28) is achieved at j , i.e.,

$$W(x) = \min_{j=1, \dots, k} \{W_j + h(x; x_j)\} = W_j + h(x; x_j).$$

Case (1). If $h(x; x_j)$ is achieved via a clockwise path, then

$$(3.31) \quad W_j + h(x; x_j) = W_j + h(x_{i+1}, x_i) + h_L(x, x_{i+1}) \geq W_j + h_L(x, x_{i+1}).$$

Case (2). If $h(x; x_j)$ is achieved via a counterclockwise path, then

$$(3.32) \quad W_j + h(x; x_j) = W_j + h(x_i, x_i) + h_L(x, x_i) \geq W_j + h_R(x, x_i).$$

Therefore, combining both cases, we have

$$(3.33) \quad W_j + h(x; x_j) \geq \min\{W_i + h_R(x; x_i), W_{i+1} + h_L(x; x_{i+1})\}.$$

From (3.18), we further know

$$(3.34) \quad \begin{aligned} W(x) &= W_j + h(x; x_j) \\ &\geq \min\{W_i + h_R(x; x_i), W_{i+1} + h_L(x; x_{i+1})\} \\ &\geq \min\{W_i + h(x; x_i), W_{i+1} + h(x; x_{i+1})\} \geq \min_{j=1, \dots, k} \{W_j + h(x; x_j)\} = W(x). \end{aligned}$$

This gives (3.30).

At last, notice that in $(x_i, x_{i+\frac{1}{2}})$, $h_R(x; x_i)$ is strictly increasing while $h_L(x; x_{i+1})$ is constant, and likewise in $(x_{i+\frac{1}{2}}, x_i)$ with $h_L(x; x_{i+1})$ being strictly decreasing and $h_R(x; x_i)$ being a constant. Thus the curves meet at most once and hence there is at most one flat connection either on the left of $x_{i+\frac{1}{2}}$ or on the right of $x_{i+\frac{1}{2}}$. This completes the proof. \square

We remark that this explicit local representation not only is helpful for constructing the global energy landscape $W(x)$ (see Figure 3) but also enables us to identify whether the least action curves track backward in time or are defined both forward and backward for the whole time $t \in \mathbb{R}$.

3.3. Consistency check for Freidlin–Wentzell variational formulas when $U(x)$ is periodic. If the original potential $U(x)$ is periodic with $\min U(x) = 0$, i.e., $\bar{b} = 0$, then $\pi_\varepsilon(x)$ is a reversible invariant measure given by $\pi_\varepsilon(x) = C_\varepsilon e^{-\frac{U(x)}{\varepsilon}}$. Since $\min U(x) = 0$, by Laplace's principle $\int e^{-\frac{U(x)}{\varepsilon}} \sim O(1)$ as $\varepsilon \rightarrow 0$. Thus $C_\varepsilon \sim O(1)$. Following Varadhan's equivalent definition [Var16, Definition 2.2] for the large deviation principle on compact domain, we compute

$$(3.35) \quad U(x) \leq \liminf_{\delta \rightarrow 0} \liminf_{\varepsilon \rightarrow 0} -\varepsilon \log \pi_\varepsilon(B_\delta(x)) \leq \limsup_{\delta \rightarrow 0} \limsup_{\varepsilon \rightarrow 0} -\varepsilon \log \pi_\varepsilon(B_\delta(x)) \leq U(x).$$

Thus $U(x)$ is the rate function for the large deviation principle of invariant measure $\pi_\varepsilon(x)$.

As a consistency check, we prove below that the constructed global energy landscape $W^*(x)$ from (3.28) and (3.29) is exactly the original potential $U(x)$.

PROPOSITION 3.4. *Let $W^*(x)$ be constructed from (3.28) and (3.29) with boundary data (3.24). If the potential $U(x)$, $x \in \mathbb{S}^1$ is periodic with $\min U(x) = 0$, then $W^*(x) = U(x)$.*

Proof. Step 1. We prove the Freidlin–Wentzell variational formula (3.22) $W_i = U(x_i) + \text{const}$.

First, define the total right/left barrier in one period as

$$(3.36) \quad J_\pm = \sum_{\ell=1}^k \left(U_{\ell \pm \frac{1}{2}} - U_\ell \right).$$

Second, for any $x_i, x_j \in \mathbb{S}^1$, by elementary calculations, we see

$$(3.37) \quad \tilde{h}_R(x_{i+1}; x_{j+1}) - \tilde{h}_R(x_i; x_{j+1}) = \begin{cases} U(x_{i+\frac{1}{2}}) - U(x_i), & j \neq i; \\ U(x_{i+\frac{1}{2}}) - U(x_i) - J_+, & j = i. \end{cases}$$

Similarly, we have

$$(3.38) \quad \tilde{h}_L(x_{i+1}; x_j) - \tilde{h}_L(x_i; x_j) = \begin{cases} -\left(U(x_{i+\frac{1}{2}}) - U(x_{i+1}) \right), & j \neq i; \\ -\left(U(x_{i+\frac{1}{2}}) - U(x_{i+1}) \right) + J_-, & j = i. \end{cases}$$

Thus we obtain

$$(3.39) \quad \begin{aligned} & \tilde{h}_R(x_{i+1}; x_{j+1}) - \tilde{h}_R(x_i; x_{j+1}) + \left(\tilde{h}_L(x_{i+1}; x_j) - \tilde{h}_L(x_i; x_j) \right) \\ &= \begin{cases} U(x_{i+1}) - U(x_i), & j \neq i; \\ U(x_{i+1}) - U(x_i) + J_- - J_+, & j = i. \end{cases} \end{aligned}$$

Third, when $U(x)$ is periodic, it is easy to verify $J_+ - J_- = 0$. Therefore,

$$(3.40) \quad \tilde{h}_R(x_{i+1}; x_{j+1}) + \tilde{h}_L(x_{i+1}; x_j) = \tilde{h}_R(x_i; x_{j+1}) + \tilde{h}_L(x_i; x_j) + U(x_{i+1}) - U(x_i).$$

Taking minimum w.r.t. j and using the definition of W_i in (3.22), we have

$$(3.41) \quad W_{i+1} = W_i + U(x_{i+1}) - U(x_i).$$

This implies

$$(3.42) \quad W_i - U(x_i) = \text{const}, \quad i = 1, \dots, k.$$

Step 2. We prove

$$(3.43) \quad U(x) = \min_{i=1, \dots, k} (U(x_i) + h(x; x_i)).$$

Fix any $x \in [i - \frac{1}{2}, i + \frac{1}{2}]$; on the one hand, from the definition of quasi-potential, we have

$$(3.44) \quad U(x) = U(x_i) + h(x; x_i).$$

On the other hand, we need to prove for any $j \neq i$

$$(3.45) \quad U(x) \leq U(x_j) + h(x; x_j).$$

From the property of $h(x; x_j)$ in Proposition 3.1, below we only prove the case that x belongs to the nonincreasing part of $h(x; x_j)$. Another case that x belongs to the nondecreasing part of $h(x; x_j)$ has the same argument.

Since x belongs to the nonincreasing part of $h(x; x_j)$ and $U(x)$ is periodic, so a clockwise path from x_j to x can be regarded as $x < x_{i+\frac{1}{2}} < x_i < \dots < x_{j-\frac{1}{2}} < x_j$. Thus

$$(3.46) \quad \begin{aligned} U(x) &\leq U(x_{i+1}) + h(x; x_{i+1}) \\ &\leq U(x_{i+2}) + h(x_{i+1}; x_{i+2}) + h(x; x_{i+1}) = U(x_{i+2}) + h(x; x_{i+2}) \\ &\leq \dots \leq U(x_j) + h(x; x_j). \end{aligned}$$

Thus, we obtain (3.43). Replace $U(x_i)$ by $U(x_i) + c = W_i$ in (3.43), and then

$$(3.47) \quad U(x) + c = \min_{i=1, \dots, k} (W_i + h(x; x_i)) = W(x). \quad \square$$

3.4. The global energy landscape $W^*(x)$ is a viscosity solution. Recalling Hamiltonian (2.5), we now prove the continuous periodic global landscape $W(x)$ constructed in (3.28) is a viscosity solution to the stationary HJE in \mathbb{S}^1 .

PROPOSITION 3.5. *Assume there are k stable local minima of $U(x)$, interleaved by k unstable local maxima indexed as (3.1) and (3.2). Let $W(x)$ be constructed in (3.28). Then*

- (i) $W(x)$ is Lipschitz continuous and periodic;
- (ii) there is at most one nondifferentiable point belonging to each increasing (resp., decreasing) interval of the original potential $U(x)$, where $W(x)$ is an increasing function connected to a constant (resp., a constant connected to a decreasing function); particularly, $W(x)$ is differentiable at all the critical points $x_i, x_{i+\frac{1}{2}}, i = 1, \dots, k$;
- (ii) $W(x)$ is a viscosity solution to HJE

$$(3.48) \quad H(W'(x), x) = W'(W' - U') = 0, \quad x \in \mathbb{S}^1,$$

and satisfies the boundary data $W(x_j) = W_j$ at $x_j, j = 1, \dots, k$.

Proof. First, from Proposition 3.1, $h(y; x_i)$ is Lipschitz continuous and periodic, so by the definition in (3.28), $W(x)$ satisfies conclusion (i).

Second, similar to the observations for $h_R(x; x_i)$ defined in (3.14), we characterize the shape of $W(x)$. For each increasing interval of $U(x)$, $h(x; x_i) + W_i$ with different x_i have only three possible shapes: constant, increasing part of $U(x) + \text{const}$, or

increasing function $U(x) + \text{const}$ connected to a constant. It is easy to verify that the minimum among all those $h(x; x_i) + W_i$ gives $W(x)$ in this increasing interval, which remains to be one of these three types. Thus there is at most one nondifferential connection point for $W(x)$, where an increasing function is connected to a constant. The scenario for each decreasing interval of $U(x)$ is similar, where the only possible connection point is a constant connected to a decreasing function. This completes conclusion (ii).

Notice the number of nondifferential points are finite and are of the same shape as in Proposition 3.1, so the verification of the viscosity solution to (3.48) of conclusion (iii) is exactly the same as that of Proposition 3.1. For the boundary conditions, recall that from Lemma 3.2, the boundary data satisfies the discrete weak KAM problem (3.24). Thus $W(x_i) = \min_{j=1, \dots, k} \{W_j + h(x; x_j)\} = W_i$ for all $i = 1, \dots, k$. This finishes the proof. \square

4. The global energy landscape $W^*(x)$ is a weak KAM solution. In this section, we first characterize the projected Aubry set \mathcal{A} and uniqueness sets for the weak KAM solutions. Then in Proposition 4.4 and Corollary 4.5, we prove the main result that the global energy landscape $W(x)$ constructed in (3.28) is a weak KAM solution, from which each calibrated curve and the projected Mather set \mathcal{M}_0 can be determined. The projected Mather set \mathcal{M}_0 is indeed the projected Aubry set \mathcal{A} , which are all the critical points of the original potential $U(x)$. Moreover, the constructed $W(x)$ is the maximal Lipschitz continuous viscosity solution satisfying the boundary data $W(x_i) = W_i$ given in (3.22). These boundary data are chosen via Lemma 3.2 so that $W^*(x) = W(x) - \min_{x \in \mathbb{S}^1} W(x)$ is the rate function for the large deviation principle of the invariant measures of the diffusion process on \mathbb{S}^1 . Hence this gives a meaningful selection principle for weak KAM solutions to (3.48). In subsections 5.4 and 5, we give more discussions on the probability interpretations and different selection principles.

4.1. Projected Aubry set \mathcal{A} and the uniqueness set. In this subsection, we first characterize the projected Aubry set \mathcal{A} , which is a uniqueness set for the weak KAM solution to HJE (3.48). We also show that all uniqueness sets must include all the local maxima/minima of $U(x)$ in Lemma 4.2. This includes the projected Mather set \mathcal{M}_0 , which is also a uniqueness set for the weak KAM solution to HJE (3.48). Second, we prove the variational formula of $W(x)$ defined in (3.28) can also be represented via the boundary data on the projected Aubry set \mathcal{A} . This representation, after being extended to the projected Aubry set \mathcal{A} , is the usually variational representation for the weak KAM solution; cf. [Tra21, Theorem 7.4]. We point out the results presented in this section are known from the general weak KAM theory. However, for our simple one-dimensional example, the proofs are explicit and simple, so we only outline the proofs for completeness.

For the one-dimensional Hamiltonian (2.5), an equivalent characterization for the projected Aubry set \mathcal{A} (4.1) is given by using the viscosity solutions to HJE (3.48).

LEMMA 4.1. *Assume there are k stable local minimums of $U(x)$, interleaved by k unstable local maximums indexed as (3.1) and (3.2). The projected Aubry set for the Hamiltonian (2.5) is*

$$(4.1) \quad \mathcal{A} := \{x_i, x_{i+\frac{1}{2}}; i = 1, \dots, k\}.$$

Proof. From [Tra21, Definition 7.32], the projected Aubry set \mathcal{A} is all the starting points x such as the Mañé potential $v(y; x)$ is a viscosity solution to (3.48) on \mathbb{S}^1 .

Indeed, from Lemma A.1 (see also [Tra21, Theorem 2.41] or [FRF09, Theorem 2.4]), we know the Mañé potential $v(y; x)$ for any $x \in \mathbb{S}^1$ is the maximal Lipschitz continuous viscosity subsolution. On the one hand, Proposition 3.1 shows that if x is a critical point, then $v(y; x) = h(y; x)$ is a viscosity solution to (3.48). Thus critical points belong to the projected Aubry set \mathcal{A} . On the other hand, for any x not being the critical points, then from Lemma A.1, the shape of $v(y; x)$ violates the viscosity supersolution test for HJE (3.48) and thus \mathcal{A} must be a subset of all the critical points. This gives the characterization of the projected Aubry set (4.1). \square

We remark there are also other characterizations for the projected Aubry set \mathcal{A} (see, for instance, [Con01, Fat08]):

$$(4.2) \quad \mathcal{A} = \{x \in \mathbb{S}^1; h(x, x) = 0\}.$$

Notice the Lagrangian $L(s, x) \geq 0$ and satisfies the property (2.7). Thus (4.2) can also be used to conclude the same characterization (4.1).

Now we recall that the projected Mather set \mathcal{M}_0 [Fat08, Theorem 4.12.6] is a uniqueness set for weak KAM solutions. Here a uniqueness set M means that if two weak KAM solutions u and \tilde{u} coincide on an M , then they must be the same everywhere. In Lemma 4.2, we prove that the projected Mather set \mathcal{M}_0 in our example must contain all the local maxima $\{x_{i+\frac{1}{2}}; i = 1, \dots, k\}$ of $U(x)$. Later in Proposition 4.4, we will prove the projected Mather set is exactly the projected Aubry set \mathcal{A} .

LEMMA 4.2. *All the uniqueness sets $M \subset \mathcal{A}$ of the weak KAM solutions to HJE (3.48) must contain all the local maxima $\{x_{i+\frac{1}{2}}; i = 1, \dots, k\}$ and local minima $\{x_i; i = 1, \dots, k\}$ of $U(x)$.*

Proof. Let $M \subset \mathcal{A}$ be a uniqueness set for the weak KAM solutions to HJE (3.48). We prove for any i the maximum point $x_{i+\frac{1}{2}} \in M$.

Using the argument by contradiction, we assume if $x_{i+\frac{1}{2}} \notin M$ for some i . Then we can choose the boundary values $W(x^*) = 0$ for all $x^* \in M$. It is easy to see zero boundary values always satisfy the consistent condition for the discrete weak KAM problem on M , i.e.,

$$(4.3) \quad W(x) = \min_{x^* \in M} \{W(x^*) + h(x; x^*)\} \quad \forall x \in M.$$

Then we use $W(x^*)$, $x^* \in M$, to construct a weak KAM solution $W(x)$, $x \in \mathbb{S}^1$. Particularly, we have

$$(4.4) \quad W(x_{i+\frac{1}{2}}) = \min_{x^* \in M} \{0 + h(x_{i+\frac{1}{2}}; x^*)\} > 0.$$

On the other hand, setting $\tilde{W}(x_{i+\frac{1}{2}}) = 0$, together with zero values in M , we can verify they still satisfy the consistent condition for the discrete weak KAM problem on the subset $M \cup \{x_{i+\frac{1}{2}}\} \subset \mathcal{A}$, and thus we can construct another weak KAM solution

$$(4.5) \quad \tilde{W}(x) = \min_{x^* \in M \cup \{x_{i+\frac{1}{2}}\}} \{W(x^*) + h(x; x^*)\} \quad \forall x \in \mathbb{S}^1.$$

One can see $\tilde{W}(x^*) = W(x^*) = 0$ for $x^* \in M$ but $\tilde{W} \neq W$ at $x_{i+\frac{1}{2}}$. This contradicts the definition of the uniqueness set. Similar arguments apply to the local minimums. \square

Remark 2. One can see an illustration of the uniqueness set in Figure 3 (right). Indeed, although $W(x_i) = 0$, $i = 1, \dots, k$, $W(x) \not\equiv 0$ in Figure 3 (right). But if we

add additional boundary values $W(x_{i+\frac{1}{2}}) = 0$ for $i = 1, \dots, k$, then the only solution is $W(x) \equiv 0$. This is because $h(y; x_{i+\frac{1}{2}}) = 0$ for $y \in [x_i, x_{i+1}]$; see Remark 1.

Next, we observe $W(x)$ in (3.28) is determined by the boundary values on the set of all the stable critical points of $U(x)$. Then the variational construction (3.28) induces the values of $W(x)$ at all the unstable critical points. After including these induced boundary values, the construction of $W(x)$ can be alternatively extended as below.

LEMMA 4.3. *Assume there are k stable local minima of $U(x)$, interleaved by k unstable local maxima periodically indexed as (3.1) and (3.2). Let $W(x)$ be defined in (3.28) and W_i be defined in (3.22). Then $W(x)$ has an alternative representation*

$$(4.6) \quad W(x) = \min_i \{W_i + h(x; x_i), W(x_{i+\frac{1}{2}}) + h(x; x_{i+\frac{1}{2}})\}.$$

Proof. Using definition (3.28), at $x_{j+\frac{1}{2}}$, we have

$$(4.7) \quad \begin{aligned} W(x_{j+\frac{1}{2}}) + h(x; x_{j+\frac{1}{2}}) &= \min_i \left(W_i + h(x_{j+\frac{1}{2}}; x_i) + h(x; x_{j+\frac{1}{2}}) \right) \\ &\geq \min_i (W_i + h(x; x_i)) = W(x), \end{aligned}$$

where we used the triangle inequality $h(z; x) + h(y; z) \geq h(y; x)$. Thus we have

$$(4.8) \quad \min_j \left(W(x_{j+\frac{1}{2}}) + h(x; x_{j+\frac{1}{2}}) \right) \geq W(x).$$

This together with $\min_i \{W_i + h(x; x_i), W(x_{i+\frac{1}{2}}) + h(x; x_{i+\frac{1}{2}})\} \leq W(x)$ implies (4.6). \square

4.2. The computation of the calibrated curves and the projected Mather set \mathcal{M}_0 . After all the preparations above, we now prove $W(x)$ constructed via (3.28) and boundary data (3.22) is a weak KAM solution of negative type.

Recall the definition of weak KAM solutions of negative type; cf. [Fat08, Definition 4.1.11]

DEFINITION 1. *We say a continuous function $u \in C(\mathbb{S}^1)$ is a weak KAM solution of negative type to HJE (3.48) if*

- (I) *u is dominated by L (denoted as $u \prec L$), i.e., for any absolutely continuous curve $\gamma \in AC([a, b]; \mathbb{S}^1)$,*

$$(4.9) \quad u(\gamma(b)) - u(\gamma(a)) \leq \int_a^b L(\dot{\gamma}, \gamma) dt;$$

- (II) *for any x , there exists a continuous, piecewise C^1 curve $\gamma: (-\infty, 0] \rightarrow \mathbb{S}^1$ with $\gamma(0) = x$ such that for any $a < b \leq 0$*

$$(4.10) \quad u(\gamma(b)) - u(\gamma(a)) = \int_a^b L(\dot{\gamma}, \gamma) dt.$$

Remark 3. Such a curve γ in condition (II) is called a calibrated curve, or a backward characteristic; see examples in Figure 4 for two calibrated curves associated with $W(x)$.

PROPOSITION 4.4. *Assume there are k stable local minimums of $U(x)$, interleaved by k unstable local maximums indexed as (3.1) and (3.2). Let $W(x)$ be defined in (3.28). Then*




FIG. 4. An illustration of the weak KAM solution $W(x)$ and two calibrated curves starting from x_1 and x_2 backward in time. The solid black line is the original potential $U(x)$ while the solid red line is the weak KAM solution $W(x)$, which serves as the global energy landscape in the zero noise limit. The calibrated curve (green arrow) starting from x_1 solves “downhill” ODE $\dot{\gamma} = -U'(\gamma)$ (4.11) backward in time and tracks back to $x_{i-\frac{1}{2}}$ with the same level in the global energy landscape $W(x)$. The ratio of the probabilities at x_1 and its reference point $x_{i-\frac{1}{2}}$ equals one, which indicates these two states appear with the same probability in the zero noise limit. The calibrated curve (orange arrow) starting from x_2 solves “uphill” ODE $\dot{\gamma} = U'(\gamma)$ (4.13) and tracks back to its reference point x_i . The probability ratio $\frac{\pi_\epsilon(x_2)}{\pi_\epsilon(x_i)}$, in the zero noise limit, is smaller than 1 and indicates state x_2 appear less likely than x_i .

- (i) for each $x \in \mathbb{S}^1$, there exists a calibrated curve tracking back to a critical point of $U(x)$;
- (ii) the projected Mather set \mathcal{M}_0 is same as the projected Aubry set $\mathcal{A} = \{x_i, x_{i+\frac{1}{2}}; i = 1, \dots, k\}$.

Proof. Recall the explicit characterization for the shape of $W(x)$. Given any $x \in \mathbb{S}^1$, we first assume x locates on a decreasing interval of $U(x)$, i.e., $x \in [x_{i-\frac{1}{2}}, x_i]$ for some i . Then from conclusion (ii) in Proposition 3.5, we know that there exists a $x^* \in [x_{i-\frac{1}{2}}, x_i]$ such that either (a) x belongs to a constant interval such that $W(x) \equiv W(x_{i-\frac{1}{2}})$ for $x_{i-\frac{1}{2}} \leq x \leq x^*$, or (b) x belongs to a decreasing interval such that $W(x) = U(x) + \text{const}$ for $x^* \leq x \leq x_i$; see Figure 4.

For case (a), we solve the following downhill ODE backward in time:

$$(4.11) \quad \dot{\gamma} = -U'(\gamma), \quad t \leq 0; \quad \gamma(0) = x.$$

Then we obtain a unique ODE solution $\gamma(t)$ with $\gamma(-\infty) = x_{i-\frac{1}{2}}$. Along this ODE solution, we verify that for any $a < b \leq 0$

$$(4.12) \quad \int_a^b L(\dot{\gamma}, \gamma) dt = \int_a^b \frac{1}{4} |\dot{\gamma} + U'(\gamma)|^2 dt = 0 = W(\gamma(b)) - W(\gamma(a)).$$

For case (b), we solve the following uphill ODE backward in time:

$$(4.13) \quad \dot{\gamma} = U'(\gamma), \quad t \leq 0; \quad \gamma(0) = x.$$

Then we obtain a unique ODE solution $\gamma(t)$ with $\gamma(-\infty) = x_i$. Along this ODE solution, we verify that for any $a < b \leq 0$

$$(4.14) \quad \int_a^b \frac{1}{4} |\dot{\gamma} + U'(\gamma)|^2 dt = \int_a^b \left(\frac{1}{4} |\dot{\gamma} - U'(\gamma)|^2 + \dot{\gamma} U'(\gamma) \right) dt \\ = U(\gamma(b)) - U(\gamma(a)) = W(\gamma(b)) - W(\gamma(a)).$$

Therefore, for both cases, we verified $W(x), x \in [x_{i-\frac{1}{2}}, x_i]$, satisfies condition (II). Similarly, if $x \in [x_i, x_{i+\frac{1}{2}}]$ for some i , one can also repeat the same argument to verify $W(x), x \in [x_i, x_{i+\frac{1}{2}}]$, satisfies condition (II).

In summary, calibrated curves have three types: Type (1) For any differential point $x \in \mathbb{S}^1$ located on a strictly increasing or decreasing part, there exists a unique backward characteristic solving (4.13) such that $\gamma(0) = x$, $\gamma(-\infty)$ tracks back to a unique local minimum (attractor) x_i in the same basin of attraction as x . Type (2) For any differential point $x \in \mathbb{S}^1$ located on a constant part of $W(x)$, there exists a unique backward characteristic solving (4.11) such that $\gamma(0) = x$, $\gamma(-\infty)$ tracks back to the local maximum $x_{i-\frac{1}{2}}$ at the end of the constant segment of $W(x)$. Type (3) For any nondifferential points $x \in \mathbb{S}^1$, there exist two backward characteristics either solving (4.13) or (4.11) and thus they track back to one of the adjacent critical points.

Consequently, based on the Aubry–Mather theory (cf. [Eva08, Fat08]), a Mather measure concentrates on one of those extremes $\gamma(-\infty)$ for the above calibrated curves and $s = 0$, i.e., $\mu = \delta(x - \gamma(-\infty))\delta(s)$. In detail, one can define μ_T for fixed T as $\langle f, \mu_T \rangle := \frac{1}{T} \int_{-T}^0 f(\dot{\gamma}(t), \gamma(t)) dt$. Then taking the limit we have

$$\int_{T\mathbb{S}^1} f(s, x) \mu(ds, dx) = \lim_{T \rightarrow +\infty} \int_{T\mathbb{S}^1} f(s, x) \mu_T(ds, dx) = \int_{T\mathbb{S}^1} f(s, x) \delta_{\dot{\gamma}(-\infty)} \otimes \delta_{\gamma(-\infty)}.$$

Thus the Mather set is given by

$$(4.15) \quad \tilde{M} = \overline{\cup \text{support } \mu} = \{(x_i, 0), (x_{i+\frac{1}{2}}, 0); i = 1, \dots, k\}.$$

Hence we conclude (i) and (ii). \square

Recall $W(x)$ is a viscosity solution to (3.48) proved in Proposition 3.5. Notice the weak KAM condition (I) can be directly implied from $W(x) \in \text{Lip}(\mathbb{S}^1)$ being a viscosity subsolution satisfying $H(W'(x), x) \leq 0$ a.e. $x \in \mathbb{S}^1$; see Proposition 3.5. Indeed, for any absolutely continuous curve $\gamma(\cdot)$ with $\gamma \in AC([a, b]; \mathbb{S}^1)$, we have

$$(4.16) \quad W(\gamma(b)) - W(\gamma(a)) = \int_a^b W'(\gamma(s)) \cdot \dot{\gamma}(s) ds \\ \leq \int_a^b (L(\dot{\gamma}(s), \gamma(s)) + H(W'(\gamma(s)), \gamma(s))) ds \leq \int_a^b L(\dot{\gamma}(s), \gamma(s)) ds.$$

The maximality of $W(x)$ is the same as Proposition 4.8, where the boundary data W_j is given only on a subset of the projected Aubry set. Thus we refer to the proof of Proposition 4.8. This, together with Proposition 4.4, yields the following.

COROLLARY 4.5. *Assume there are k stable local minimums of $U(x)$, interleaved by k unstable local maximums indexed as (3.1) and (3.2). Let $W(x)$ be defined in (3.28). Then $W(x)$ is a weak KAM solution of negative type to HJE (3.48). And $W(x)$ is the maximal Lipschitz continuous viscosity solution to (3.48) that satisfy boundary data $W(x_j) = W_j$ at x_j , $j = 1, \dots, k$.*

We point out that we only need the boundary data on the local minima of U , i.e., a subset of the projected Aubry set, to select a meaningful weak KAM solution which serves as the rate function in the large deviation principle for invariant measures. On the other hand, given any boundary data in a subset of the projected Aubry set, one can construct a maximal Lipschitz continuous viscosity solution; see Proposition 4.8.

Remark 4. From (4.12), it is easy to see that for any curve $\gamma : \mathbb{R} \rightarrow \mathbb{S}^1$ which solves the ODE $\dot{\gamma} = \pm U'(\gamma), t \in \mathbb{R}$, is a least action curve. This gives the projected Mañé set

$$\overline{\cup\{\gamma(t); \gamma \text{ solves } \dot{\gamma} = \pm U'(\gamma), t \in \mathbb{R}\}} = \mathbb{S}^1.$$

The Mañé set itself is the collection of the Lagrangian graph $(\gamma, \pm U(\gamma))$ of those least action curves. Furthermore, in this simple example, it is easy to see the Mather set (4.15) is a compact Lipschitz graph and is invariant under the Euler–Lagrange flow. This is the essence of the celebrated Mather graph theorem that characterizes the graph property of the Mather set.

4.2.1. Invariant solutions of the Lax–Oleinik semigroup. In this section, using the equivalent characterization of invariant solutions of the Lax–Oleinik semigroup [Fat08, Proposition 4.6.7], we give a direct corollary that $W(x)$ defined in (3.28) is an invariant solution for the Lax–Oleinik semigroup S_t associated with the dynamic HJE $\partial_t u + H(\partial_x u(x), x) = 0$, i.e., for $t \geq 0$,

$$(4.17) \quad (S_t u_T)(y) := \inf_x \left(u_T(x) + \inf_{\gamma; \gamma(0)=x, \gamma(t)=y} \int_0^t L(\dot{\gamma}, \gamma) d\tau \right) \quad \forall u_T(x).$$

COROLLARY 4.6. *Any weak KAM solution $w(x)$ to HJE (3.48) is an invariant solution of the Lax–Oleinik semigroup S_t and satisfies the representation*

$$(4.18) \quad w(y) = \inf_{x \in \mathbb{S}^1} (w(x) + v(y; x)) = \inf_{x_i \in \mathcal{A}} (w(x_i) + h(y; x_i)).$$

Particularly, $W(x)$ defined in (3.28) is an invariant solution of the Lax–Oleinik semigroup S_t .

Proof. First, from [Fat08, Proposition 4.6.7], any weak KAM solution $w(x)$ is an invariant solution of the Lax–Oleinik semigroup S_t . Thus

$$w(y) = \inf_x \left(w(x) + \inf_{\gamma; \gamma(0)=x, \gamma(t)=y} \int_0^t L(\dot{\gamma}, \gamma) d\tau \right).$$

Taking infimum w.r.t. t and exchanging \inf_x and \inf_t , we obtain $w(y) = \inf_x (w(x) + v(y; x))$.

Second, take the boundary values $w(x_i)$ on the projected Aubry set \mathcal{A} . Since the projected Aubry set is a uniqueness set for the weak KAM solution [Fat08, Theorem 4.12.6], these boundary values can uniquely define a weak KAM solution. Meanwhile, from Corollary 4.5 $w(x) = \inf_{x_i \in \mathcal{A}} (w(x_i) + h(y; x_i))$ is a weak KAM solution and thus the representation (4.18) holds uniquely. \square

Remark 5. We remark that for a compact domain, the existence of invariant solutions of S_t and the convergence from a dynamic solution to an invariant solution were proved in [Fat08, NR99]. However, the invariant solutions are not unique. The weak KAM solutions are also not unique; see examples in Figure 3 and [FRF09].

4.3. Generating a set of consistent boundary data and constructing a maximal Lipschitz continuous viscosity solution. In this subsection, given any nonconsistent boundary data on a subset of the projected Aubry set, we can first use it to generate a set of consistent data satisfying the discrete weak KAM problem. Then based on these consistent data, we prove the variational formula $W(x)$ is the maximal Lipschitz continuous viscosity solution to the HJE satisfying the generated boundary data.

4.3.1. Nonconsistent boundary data induce a set of consistent data satisfying (3.24). Now given any boundary data

$$(4.19) \quad \{W_\ell\} \quad \text{at } D := \{x'_\ell; \ell = 1, 2, \dots, m\} \subset \mathcal{A},$$

which may not satisfy the discrete weak KAM problem (3.24), the following procedure can be used to obtain a set of consistent boundary data satisfying (3.24). For any $j = 1, \dots, m$, define

$$(4.20) \quad W(x) = \min_{\ell=1, \dots, m} \{W_\ell + h(x; x'_\ell)\}.$$

Then $\tilde{W}_\ell := W(x'_\ell)$ is a set of consistent data satisfying discrete weak KAM problem

$$(4.21) \quad \tilde{W}_j = \min_{\ell=1, \dots, m} \{\tilde{W}_\ell + h(x'_j; x'_\ell)\}, \quad j = 1, \dots, m.$$

PROPOSITION 4.7. *Given any boundary data $\{W_\ell\}$ on a subset $\{x'_\ell; \ell = 1, 2, \dots, m\}$ of \mathcal{A} , then*

$$(4.22) \quad \tilde{W}_j := \min_{\ell=1, \dots, m} \{W_\ell + h(x'_j; x'_\ell)\}, \quad j = 1, \dots, m,$$

satisfy the discrete weak KAM problem (4.21).

Proof. Given boundary data $\{W_\ell\}$, $\ell = 1, \dots, m$, let $W(x)$ be defined in (4.20). On the one hand, $\tilde{W}_j := W(x'_j) \leq W_j$ for any $j = 1, \dots, m$. Thus

$$(4.23) \quad \tilde{W}(x) = \min_{\ell=1, \dots, m} \{\tilde{W}_\ell + h(x; x'_\ell)\} \leq W(x).$$

On the other hand, from Proposition 3.5, $W(x)$ is a Lipschitz continuous viscosity solution, while from Proposition 3.1, $\tilde{W}_\ell + h(x; x'_\ell)$ is the maximal Lipschitz continuous viscosity solution. Thus from $W(x'_\ell) = \tilde{W}_\ell$, we have

$$(4.24) \quad W(x) \leq \min_{\ell=1, \dots, m} \{\tilde{W}_\ell + h(x; x'_\ell)\} = \tilde{W}(x).$$

Therefore, we conclude $\tilde{W}(x) = W(x)$. Particularly, $\tilde{W}(x'_\ell) = W(x'_\ell) = \tilde{W}_\ell$ for $\ell = 1, \dots, m$ and thus \tilde{W}_ℓ is a consistent boundary value satisfying (4.21). \square

We point out in the above proposition that the subset $\{x'_\ell; \ell = 1, 2, \dots, m\}$ is not necessarily a uniqueness set. Indeed, in the next subsection, we will prove that as long as the boundary values W_ℓ satisfy the discrete weak KAM problem (4.21), then we can use those data to obtain a maximal Lipschitz continuous viscosity solution. Particularly, the weak KAM solution in Corollary 4.5 is the maximal Lipschitz continuous viscosity solution satisfying boundary data (3.22).

4.3.2. Maximal Lipschitz continuous viscosity solution based on consistent data. In the next proposition, given any boundary values W_ℓ , for $D := \{x'_\ell; \ell = 1, \dots, m\} \subset \mathcal{A}$, if $\{W_\ell; \ell = 1, \dots, m\}$ satisfies the discrete weak KAM problem (4.21), we prove $W(x)$ with the representation

$$(4.25) \quad W(x) = \min_{\ell=1, \dots, m} \{W_\ell + h(x; x'_\ell)\}$$

is indeed the maximal Lipschitz continuous viscosity solution to the HJE satisfying given boundary values W_ℓ , i.e.,

$$(4.26) \quad H(W'(x), x) = W'(W' - U') = 0, \quad x \in \mathbb{S}^1; \quad W(x'_\ell) = W_\ell \text{ for } x'_\ell \in D.$$

Consequently, $W(x)$ constructed in (3.28) is not only one of the weak KAM solution satisfying a given boundary condition on all local minimums but also the maximal Lipschitz continuous viscosity solution to $H(W'(x), x) = W'(W' - U') = 0$ with those given boundary conditions.

PROPOSITION 4.8. *Given any boundary values W_ℓ on $D = \{x'_\ell; \ell = 1, \dots, m\} \subset \mathcal{A}$, the solution $W(x)$ constructed via (4.25) is the maximal Lipschitz continuous viscosity solution to (4.26).*

Proof. From Step 4 in the proof of Proposition 3.1, we know that for any $\ell = 1, \dots, m$, the lifted Peierls barrier $W_\ell + h(x; x'_\ell)$ is the maximal Lipschitz continuous viscosity solution to $W'(W' - U') = 0$ satisfying the boundary value $W(x'_\ell) = W_\ell$. Given any Lipschitz continuous viscosity solution $\tilde{W}(x)$ to (4.26), since $\tilde{W}(x)$ is a Lipschitz continuous viscosity subsolution, we know

$$(4.27) \quad \tilde{W}(x) \leq W_\ell + h(x; x'_\ell)$$

for any $x'_\ell \in D$. Notice $\tilde{W}(x)$ satisfies all the boundary values $\tilde{W}(x'_\ell) = W_\ell$ for $x'_\ell \in D$, hence taking the minimum for all $x'_\ell \in D$, we obtain

$$(4.28) \quad \tilde{W}(x) \leq \min_{\ell=1, \dots, m} \{W_\ell + h(x; x'_\ell)\} = W(x).$$

Thus, together with Proposition 3.5, $W(x)$ is the maximal Lipschitz continuous viscosity solution satisfying all the boundary values $W(x'_\ell) = W_\ell$ on D . \square

5. Selection principles for weak KAM solutions and probability interpretation. In this section, we discuss the selection principle given by the global energy landscape in the large deviation principle for invariant measures and also compare it with another selection principle via the vanishing discount limit. It is well known that the dynamic HJE has a unique viscosity solution and the long time limits exist but are not unique. That is to say, a selection principle is needed for viscosity solutions to stationary HJE, even when the Hamiltonian is strictly convex. Among all the viscosity solutions, the weak KAM solutions are the maximal Lipschitz continuous viscosity solutions satisfying specific boundary conditions on a uniqueness set. Hence a weak KAM solution serves as a natural candidate for the selection principle and the key point to select a meaningful weak KAM solution is the determination of boundary data on a uniqueness set of the weak KAM solutions. The variational formula for those boundary data $W(x_i)$ obtained in (3.22) gives a unique determination that captures the asymptotic behaviors of the original stochastic process at each local attractor. Equivalently, we summarize it as a selection principle for those weak KAM solutions by exchanging the double limits for t and ε . That is, we first take the

long time limit $\lim_{t \rightarrow +\infty} \rho_\varepsilon(x, t)$ which is unique due to ergodicity and then take the zero noise limit $\varepsilon \rightarrow 0$ due to the large deviation principle for invariant measures. In general, the vanishing viscosity limit is an approximation method for stationary HJE, but the limit is only in the subsequence sense and is not unique. Our selection principle provides a special viscosity approximation to the stationary HJE whose vanishing viscosity limit is unique. The probability interpretations via the Boltzmann analysis of the global energy landscape from the weak KAM perspective are also discussed.

5.1. Examples for nonuniqueness. In the proof of Corollary 4.5, we did not use the explicit values of $W(x_i)$ at the local minima x_i . Indeed, given any boundary values $W(x_i)$ for any subset of (not necessarily all) those local minima x_i , as long as those boundary values are consistent with the associated discrete weak KAM problem (3.24), then $W(x)$ determined by those given boundary values though (3.28) is a weak KAM solution.

Furthermore, we use a classical example, which appeared in the first edition of the book [FW12, section 6.4] in 1979, to illustrate the boundary values consistent with the discrete weak KAM problem (3.24) are not unique.

Choose a skew periodic potential $U(x)$ such that U has three local minima x_1, x_2, x_3 with values 1, 0, 2 and has four local maxima $x_{\frac{1}{2}}, x_{1+\frac{1}{2}}, x_{2+\frac{1}{2}}, x_{3+\frac{1}{2}}$ with values 7, 5, 10, 11. In Figure 3, the original skew periodic potential $U(x)$ is the same as the one in Figure 5, and two plots for $W^*(x)$ with different boundary values based on (5.1) and (5.2) are shown for comparison.

One has a set of boundary values computed from (3.22), $W(x_1) = 13, W(x_2) = 12, W(x_3) = 11$. It is easy to verify that this set of boundary values satisfies the discrete weak KAM problem (3.24). Then from (3.28) and (3.29), $W(x)$ is given by




FIG. 5. The comparison between the Peierls barrier $h(y; x_2)$ and a general Mañé potential $v(y; x^{**})$. The dashed black line is the skew periodic potential $U(x)$ with three local minimums 1, 0, 2 in $\mathbb{S}^1 = [6L, 6]$. The solid green line is a Peierls barrier $h(y; x_2)$ starting from a local minimum $x_2 = 3$, which is periodic with one constant-cut from above and only one nondifferentiable point x^* . The left slope is larger than the right slope at x^* , so $h(x^*; x_2)$ satisfies the viscosity solution test. The solid purple line starting from x^{**} is the left half of the Mañé potential $v(y; x^{**})$, which has an additional constant-cut from below at the nondifferentiable point x^{**} . The left slope is smaller than the right slope at x^{**} , so $v(x^{**}; x^{**})$ violates the viscosity supersolution test (i.e., violates the entropy condition).

$$(5.1) \quad W^*(x) = W(x) - \min_i W(x_i) = \begin{cases} 1 + U(x), & x \in [x_{\frac{1}{2}}, x_2]; \\ \min\{1 + U(x), 8\}, & x \in [x_2, x_{2+\frac{1}{2}}]; \\ U(x) - 2, & x \in [x_{2+\frac{1}{2}}, x_3]; \\ \min\{U(x) - 2, 8\}, & x \in [x_3, x_{3+\frac{1}{2}}], \end{cases}$$

which satisfies Proposition 3.5.

Another set of boundary values can be chosen as $W(x_1) = W(x_2) = W(x_3) = 0$. It is easy to verify that this set of boundary values also satisfy the discrete weak KAM problem (3.24). Then from (3.28), $W(x)$ is given by

$$(5.2) \quad W^*(x) = W(x) = \begin{cases} U(x) - 1, & x \in [x_{\frac{1}{2}}, x_{1+\frac{1}{2}}]; \\ \min\{U(x), 4\}, & x \in [x_{1+\frac{1}{2}}, x_2]; \\ \min\{U(x), 8\}, & x \in [x_2, x_{2+\frac{1}{2}}]; \\ U(x) - 2, & x \in [x_{2+\frac{1}{2}}, x_3]; \\ \min\{U(x) - 2, 6\}, & x \in [x_3, x_{3+\frac{1}{2}}]. \end{cases}$$

which also satisfies Proposition 3.5. From Corollary 4.5, both sets of boundary values give a weak KAM solution to (3.48), so weak KAM solutions are not unique.

5.2. Exchange limits $\varepsilon \rightarrow 0$, $t \rightarrow +\infty$ in two large deviation principles.

Below we discuss a special case for which the long time behavior limit $t \rightarrow +\infty$ and the zero noise limit $\varepsilon \rightarrow 0$ for the diffusion process (2.1) can be exchanged. Notice that in general, it is not exchangeable.

Recall the Fokker–Planck equation on \mathbb{S}^1 (2.3) and WKB reformulation $\rho_\varepsilon(x, t) = e^{-\frac{\psi_\varepsilon(x, t)}{\varepsilon}}$. The viscous HJE associated with ψ_ε is

$$(5.3) \quad \partial_t \psi_\varepsilon(x, t) + H(\partial_x \psi_\varepsilon(x, t), x) = \varepsilon (\partial_{xx} \psi_\varepsilon(x, t) - U''(x)), \quad x \in \mathbb{S}^1, t > 0.$$

In general, the two limits for $\psi_\varepsilon(x, t)$ as $\varepsilon \rightarrow 0$ and $t \rightarrow +\infty$ cannot be exchanged. But with special initial data, we have the following result.

PROPOSITION 5.1. *Let $W^*(x)$ be the rate function given by (3.29) and assume $U(x)$ have finite local extremes. Assume $\rho_\varepsilon(x, t)$ is a solution to the Fokker–Planck equation (2.3) and the initial distribution is given by $\rho_\varepsilon(x, 0) = C_\varepsilon e^{-\frac{W^*(x)}{\varepsilon}}$. Then*

$$(5.4) \quad \lim_{\varepsilon \rightarrow 0} \left(-\varepsilon \log \lim_{t \rightarrow +\infty} \rho_\varepsilon(x, t) \right) \text{ “=” } W^*(x) = \lim_{t \rightarrow +\infty} \lim_{\varepsilon \rightarrow 0} -\varepsilon \log \rho_\varepsilon(x, t).$$

Here “=” is in the large deviation principle sense (2.20).

Proof. On the one hand, for fixed ε , the ergodicity $\lim_{t \rightarrow +\infty} \rho_\varepsilon(x, t) = \pi_\varepsilon(x)$ is a standard result for overdamped Langevin dynamics on \mathbb{S}^1 . Thus from the large deviation principle (2.20) [FW12, Chapter 6, Theorem 4.3], we have

$$(5.5) \quad \lim_{\varepsilon \rightarrow 0} \left(-\varepsilon \log \lim_{t \rightarrow +\infty} \rho_\varepsilon(x, t) \right) \text{ “=” } W^*(x),$$

where “=” is in the large deviation principle sense (2.20).

On the other hand, $\psi_\varepsilon(x, t) = -\varepsilon \log \rho_\varepsilon(x, t)$ is the solution to the HJE (5.3) with initial data $W^*(x) - \varepsilon \log C_\varepsilon$. Since $\min W^*(x) = 0$, by the Laplace principle $\int e^{-\frac{W^*(x)}{\varepsilon}} \sim O(1)$ as $\varepsilon \rightarrow 0$. Thus $C_\varepsilon \sim O(1)$ and as $\varepsilon \rightarrow 0$, $W^*(x) - \varepsilon \log C_\varepsilon \rightarrow W^*(x)$. From the [CL83, CL84] vanishing viscosity method, we know the convergence from

the solution $\psi_\varepsilon(x, t)$ of (5.3) to the viscosity solution $\psi(x, t)$ of the limiting first order HJE

$$\partial_t \psi(x, t) + H(\partial_x \psi(x, t), x) = 0, \quad \psi(x, 0) = W^*(x).$$

Then by the Lax–Oleinik semigroup representation

$$(5.6) \quad \psi(x, t) = (S_t W^*)(x) = W^*(x),$$

where we used that $W^*(x)$ is an invariant solution due to Corollary 4.6. Thus we know

$$W^*(x) = \lim_{t \rightarrow +\infty} W^*(x) = \lim_{t \rightarrow +\infty} \lim_{\varepsilon \rightarrow 0} -\varepsilon \log \rho_\varepsilon(x, t). \quad \square$$

We remark that the exchanging of two limits on the left- and right-hand sides is in general incorrect. Indeed, the limits in the left-hand side of (5.4) are unique. This is because the invariant measure for $t \rightarrow +\infty$ exists and is unique. Then the rate function of the large deviation principle for invariant measures is unique. However, the right-hand side first finds the rate function for the large deviation principle as $\varepsilon \rightarrow 0$ at finite time, which solves a dynamic HJE. Then the long time limit $t \rightarrow +\infty$ for the dynamic solution exists [BS00] but in general is not unique. Therefore, a selection principle is needed, and particularly the limits on the left-hand side provide a meaningful selection principle for stationary HJE via the large deviation principle for invariant measures. Below, we discuss two selection principles: large deviation principle versus vanishing discount limit.

5.3. Two selection principles: Large deviation principle versus vanishing discount limit. A selection principle gives a meaningful principle to determine boundary values on the projected Aubry set \mathcal{A} . The global energy landscape $W(x)$ in (3.28), particularly the globally adjusted boundary values on the local minima (3.22), is constructed so that $W(x)$ is the rate function for the large deviation principle of the invariant measure for the diffusion process on \mathbb{S}^1 [FW12, Chapter 6, Theorem 4.3]. That is to say, the large deviation rate function $W(x)$ for the diffusion process serves as a selection principle for weak KAM solutions. This selection principle could also apply to other Hamiltonian dynamics with an underlying stochastic process and a large deviation principle. We formally describe this framework below for a chemical reaction process with a random-time changed Poisson representation; cf. [AK15, GL22]. For any fixed time $0 < t < +\infty$, the large deviation principle for the chemical reaction process in the thermodynamic limit was proved in [GL23] by using the convergence from the Varadhan’s discrete nonlinear semigroup $w_\varepsilon(x, t)$ to the viscosity solution $w(x, t)$ of the dynamic HJE, which has a Lax–Oleinik semigroup representation. If this Lax–Oleinik semigroup has an invariant solution, denoted as $w(x)$, then this invariant solution is a weak KAM solution and has the representation $w(x) = \inf_{y \in \mathbb{R}^d} (w(y) + v(x; y))$ via the Mañé potential $v(x; y)$; see [Tra21, Proposition 6.11, Theorem 7.5], [Fat08, Proposition 4.6.7] for proofs for a periodic domain. Notice these invariant solutions are in general not unique. However, the Lagrangian $L(s, x)$ in the least action problem $v(x; y)$ is always nonnegative and it is proved in [GL23] that the zero-cost flow (a.k.a. the dynamics following the law of large numbers) is given by $\dot{x} = \partial_p H(0, x)$. Thus the projected Aubry set \mathcal{A} , which is assumed to contain only finite many points, can be characterized by using the roots of $\partial_p H(0, x) = 0$. Then the weak KAM representation can be reduced to $w(x) = \inf_{y \in \mathcal{A}} (w(y) + h(x; y))$

[Tra21, Theorem 7.40]. Assume furthermore $w(x), x \in \mathcal{A}$, is chosen such that $w(x)$ is the rate function for the invariant measure of the chemical reaction process; then this gives a selection principle to those weak KAM solutions.

We comment on the stationary HJE (5.3) for ψ_ε with viscosity terms can be one nontrivial viscosity approximation for the stationary HJE. In general, we know the nonuniqueness for the vanishing viscosity limit of stationary HJE. How to construct a vanishing viscosity approximation to stationary HJE which has a unique limit for all vanishing ε is still an open question [Tra21]. In our example, thanks to the inhomogeneous term $\varepsilon U''(x)$ in

$$(5.7) \quad H(\partial_x \psi_\varepsilon(x, t), x) = \varepsilon (\partial_{xx} \psi_\varepsilon(x, t) - U''(x)),$$

one has a nontrivial solution but also has a uniform limit as $\varepsilon \rightarrow 0$; see (5.5). This serves as a meaningful vanishing viscosity approximation but in general, we do not have an answer.

In another direction, a selection principle is given by choosing the boundary values on the projected Aubry set \mathcal{A} so that the weak KAM solution $W(x)$ is the unique viscosity solution which is the vanishing discount limit of the solution ψ_λ to $\lambda \psi_\lambda + H(\nabla \psi_\lambda(x), x) = 0$ as $\lambda \rightarrow 0$. This direction has been widely studied in both compact and noncompact domains [Con01, Gom08, DFIZ16, IS20]. We refer to [CGMT15, MT17], which include a degenerate diffusion term in the vanishing discount limit problem, and to [IMT17] for a duality framework in the vanishing discount problem for fully nonlinear, degenerate elliptic Hamiltonian. The vanishing discount limit method is different from the vanishing viscosity limit we constructed. Particularly, for our one-dimensional example on \mathbb{S}^1 , the vanishing discount limit of the discounted HJE with the same Hamiltonian

$$(5.8) \quad \lambda \psi_\lambda + \psi'_\lambda (\psi'_\lambda - U') = 0$$

is trivial. For $\lambda > 0$, there is a unique viscosity solution $\psi_\lambda \equiv 0$ due to the comparison principle. Thus its vanishing discount limit $\lim_{\lambda \rightarrow 0^+} \psi_\lambda \equiv 0$ is the selected weak KAM solution to (3.48) via the vanishing discount limit.

Based on the discussions above, we can see at least for the diffusion process on \mathbb{S}^1 , the vanishing discount limit and the rate function in the large deviation principle are two different selection principles which result to different weak KAM solutions. This is analogous to the idea that in general the two limits $t \rightarrow +\infty$ and $\varepsilon \rightarrow 0$ for (5.3) are nonexchangeable.

5.4. Boltzmann analysis for the weak KAM solution $W^*(x)$ selected via large deviation principle. In this section, based on the weak KAM solution $W^*(x)$ defined in (3.28) with boundary data $W(x_i) = W_i$ constructed in (3.22), we elaborate some probability interpretations that can be explained or computed via the weak KAM solution properties.

The classical Boltzmann analysis in statistical mechanics shows that in an equilibrium system, the probability for a particle being at a certain state x is a function of the state's energy $E(x)$ and the temperature T ,

$$(5.9) \quad \pi(x) \propto e^{-\frac{E(x)}{k_B T}}.$$

Then the ratio of the probability between any two states is

$$(5.10) \quad \frac{\pi(x_1)}{\pi(x_2)} = e^{\frac{E(x_2) - E(x_1)}{k_B T}}.$$

However, for a nonequilibrium system, for instance, the irreversible diffusion example on \mathbb{S}^1 (2.8), this ratio cannot be computed directly from the original potential energy $U(x)$.

Indeed, the weak KAM solution $W(x)$ provides the answer, which not only serves as the good rate function of the large deviation principle of invariant measure $\pi_\varepsilon(x)$ but also allows one to find a calibrated curve for any x tracking back to a critical point in the projected Aubry set \mathcal{A} . This calibrated curve allows one to compute the ratio of the probabilities between the starting point $\gamma(0) = x^*$ and its reference point $\gamma(-\infty)$,

$$(5.11) \quad \frac{\pi_\varepsilon(x^*)}{\pi_\varepsilon(x(-\infty))} \approx e^{\frac{W(x(-\infty)) - W(x^*)}{\varepsilon}}.$$

The value of this ratio, depending on the explicit calibrated curve starting from x^* , is either 1 or $e^{\frac{U(x_i) - U(x^*)}{\varepsilon}}$. These ratios of probabilities w.r.t. different reference points due to different calibrated curves are shown in Figure 4.

Appendix A. Remarks on Mañé potential not being a viscosity solution on \mathbb{S}^1 . Regarding the conclusions (iii) and (iv) in Proposition 3.1, we emphasize that the nondifferential point cannot result from a C^1 function cut off from below by a constant; otherwise it is not a viscosity solution to HJE. Indeed, from the proof of (iv), if a C^1 function is cut off from below by a constant, then at the nondifferential point, a constant is connected to an increasing function, where $D^+h(x^*) = \emptyset$, $D^-h(x^*) = \{q; 0 \leq q \leq h'(x_+^*) = U'(x^*)\}$. Then it's easy to verify $q(q - U'(x)) \leq 0$ does not satisfy the viscosity supersolution test; see Figure 5 for the comparison of the shape of the Peierls barrier $h(y; x_i)$ and a general Mañé potential $v(y; x^{**})$.

As a byproduct, we also characterize the shape of the Mañé potential $v(y; x^{**})$ and explain why we do not use the Mañé potential to construct a global energy landscape.

LEMMA A.1. *Let the Mañé potential $v(y; x^{**})$ be defined in (3.4) and assume x^{**} is not a critical point of $U(x)$.*

- (i) *The Mañé potential $v(y; x^{**})$ is Lipschitz continuous and periodic.*
- (ii) *The starting point x^{**} must be a nondifferential point where either a constant is connected to an increasing function or a decreasing function is connected to a constant. That is to say, $v(y; x^{**})$ is a C^1 function cut off at least once by a constant zero from below.*
- (iii) *Another possible nondifferential point is same as that for the Peierls barrier.*
- (iv) *$v(y; x_i)$ is the maximal Lipschitz continuous viscosity subsolution to HJE,*

$$(A.1) \quad H(v'(y), y) = v'(v' - U') = 0, \quad y \in \mathbb{S}^1,$$

*satisfying $v(x^{**}; x^{**}) = 0$, but it does not satisfy the viscosity supersolution test at x^{**} . In other words, $\rho(y) := v'(y; x^{**})$ is not a stationary entropy shock at x^{**} to the corresponding Burgers transport equation*

$$(A.2) \quad \partial_t \rho + \partial_y(\rho^2) - \partial_y(U' \rho) = 0.$$

Proof. First, we consider the i th well of $U(x)$ containing the starting point x^{**} . Assume $x^{**} \in (x_i, x_{i+\frac{1}{2}})$, then

$$(A.3) \quad v(y; x^{**}) := \begin{cases} U(y) - U(x_i), & y \in (x_{i-\frac{1}{2}}, x_i); \\ \max\{U(y) - U(x^{**}), 0\}, & y \in (x_i, x_{i+\frac{1}{2}}). \end{cases}$$

This means at x^{**} , a constant 0 is connected to an increasing function $U(y) - U(x^{**})$. Similarly, if $x^{**} \in (x_{i-\frac{1}{2}}, x_i)$, we obtain at x^{**} , a decreasing function $U(y) - U(x^{**})$ is connected to a constant. This proves conclusion (ii).

Second, for y outside i th well, the construction is the same as the Peierls barrier. Thus conclusions (i) and (iii) follow.

Third, we only need to verify the viscosity solution test at the nondifferential point x^{**} . If the nondifferential point is a constant 0 connected to an increasing function $U(y) - U(x^{**})$, then $D^+v(x^{**}) = \emptyset$, $D^-v(x^{**}) = \{q; 0 \leq q \leq v'(x_+^{**}) = U'(x^{**})\}$. Then it's easy to verify $q(q - U'(x)) \leq 0$ satisfies the subsolution condition but does not satisfy the viscosity supersolution condition. Again, from Step 4 in the proof of Proposition 3.1, we know the Mañé potential $v(y; x^{**})$ is the maximal Lipschitz continuous viscosity subsolution to (A.1).

Last, we take $\rho(y) = v'(y; x^{**})$, and then the solution to (A.1) is equivalent to the stationary solution $\rho(y)$ to Burgers' transport equation (A.2). The stationary shock solution $\rho(y)$ at the jump point x^{**} , with the left limit ρ_- and right limit ρ_+ , satisfies

$$\rho_+^2 - U'(x^{**})\rho_+ = \rho_-^2 - U'(x^{**})\rho_-.$$

The entropy condition for a shock solution is that for any convex entropy function $\eta(\rho)$,

$$(A.4) \quad \partial_t \eta(\rho) + \partial_y (\Phi(\rho) - U' \eta(\rho)) + (\eta(\rho) - \rho \eta'(\rho)) U'' \leq 0$$

in the distribution sense. Here $\Phi'(\rho) = 2\rho \eta'(\rho)$. For scalar equations, one can just take $\eta(\rho) = \rho^2$ and thus $\Phi(\rho) = \frac{4}{3}\rho^3$. Then the entropy condition (A.4) for the stationary shock $\rho(y)$ at x^{**} becomes

$$(A.5) \quad \Phi(\rho_+) - U'(x^{**})\eta(\rho_+) - (\Phi(\rho_-) - U'(x^{**})\eta(\rho_-)) = \frac{1}{3}(\rho_+ - \rho_-)^3 \leq 0,$$

which implies $\rho(y)$ only has jump discontinuity at x^{**} and the left limit ρ_- is larger than the right limit ρ_+ . Back to $v(y; x^{**})$, the entropy condition is violated at x^{**} since $v'(x_+^{**}; x^{**}) > v'(x_-^{**}; x^{**})$. This entropy condition violation argument is equivalent to the violation of the viscosity supersolution condition for the Mañé potential $v(y; x^{**})$. \square

Appendix B. Remark on the weak KAM solutions of positive type.

One can also define a weak KAM solution of positive type; the only difference in the theory is a time direction. That is to say, the calibrated curve is defined on $[0, +\infty)$ and for any $0 \leq a < b$ the least action is achieved,

$$(B.1) \quad u_+(\gamma(b)) - u_+(\gamma(a)) = \int_a^b L(\dot{\gamma}, \gamma) dt.$$

Moreover, the weak KAM solution of positive type u_+ can be equivalently characterized as a invariant solution to the Lax–Oleinik semigroup S_t^+ associated with the dynamic HJE [Eva08],

$$(B.2) \quad \partial_t u - H(\partial_x u(x), x) = 0, \quad x \in \mathbb{S}^1, \quad u(x, 0) = u_0.$$

The viscosity solution to (B.2) is represented as the backward semigroup S_t^+ , i.e., for $t \geq 0$,

$$(B.3) \quad (S_t^+ u_0)(y) := \sup_x \left(u_0(x) - \inf_{\gamma; \gamma(0)=y, \gamma(t)=x} \int_0^t L(\dot{\gamma}, \gamma) d\tau \right), \quad y \in \mathbb{S}^1.$$

Then u_+ is the invariant solution of S_y^+ satisfying

$$(B.4) \quad u_+(y) = (S_t^+ u_+)(y)$$

and it is a viscosity solution to the stationary HJE [Eva08, Theorem 3.1],

$$(B.5) \quad -H(\partial_x u_+(x), x) = 0, \quad x \in \mathbb{S}^1.$$

It worth noting that the weak KAM solution of positive type $u_+(x)$ is *not* the same as the negative ones in general. But the weak KAM solution of positive type $u_+(x)$ can be constructed via the negative type ones with a time reversed Hamiltonian. Precisely, define the time reversed Hamiltonian as $\hat{H}(p, x) = H(-p, x)$ and the corresponding time reversed Lagrangian is $\hat{L}(s, x) = L(-s, x)$. Then it is easy to see the weak KAM solution of negative type, denoted as $\hat{u}_-(x)$, for the HJE

$$\hat{H}(\partial_x \hat{u}_-(x), (x)) = 0$$

satisfies the relation

$$(B.6) \quad -\hat{u}_-(x) = u_+(x),$$

where $u_+(x)$ is a weak KAM solution of positive type for the HJE,

$$-H(\partial_x u_+(x), (x)) = 0.$$

Apparently, at nondifferential points, the viscosity solution test is different for the above two stationary HJEs. For instance, in our \mathbb{S}^1 Langevin dynamics example, $\hat{H}(p, x) = p(p + U')$, and thus by Proposition 3.3, $\hat{u}_-(x)$ can be expressed as the local trimming from above of $-U$, i.e.,

$$(B.7) \quad \hat{u}_-(x) = \min\{-U, \text{const}\} \quad \text{for } x \in (x_{i-\frac{1}{2}}, x_i) \text{ or } x \in (x_i, x_{i+\frac{1}{2}}).$$

In terms of u_+ , this is a local trimming from below of U , i.e.,

$$(B.8) \quad u_+(x) = -\hat{u}_-(x) = -\min\{-U, \text{const}\} = \max\{U, -\text{const}\}.$$

However, when the potential U is periodic, i.e., the Langevin dynamics is reversible, no cutoff from above/below is performed, and thus the positive type weak KAM solution $u_+(x)$ given by (B.6) is same as the negative type $u_-(x)$ constructed via the large deviation principle (see Corollary 4.5). Indeed, $\hat{H}(p, x) = p(p + U')$ and $\hat{u}_-(x) = -U(x)$ is a weak KAM solution of the negative type associated with $\hat{H}(\partial_x \hat{u}_-(x), (x)) = 0$, which is actually solved in the classical sense. Thus $u_+(x) = U(x) = u_-(x)$. This argument is no longer true for the irreversible process, i.e., $U(x)$ is not periodic.

Appendix C. Freidlin–Wentzell variational formula. In this section, we give a coarse grained Markov chain interpretation for Freidlin–Wentzell variational formula (3.28).

To study the multiwell exit problem, the essential idea follows Kolmogorov's construction of a Markov chain induced by the continuous process X_t in (2.1). Denote the collection of all the local minimums as $\Gamma := \{x_i, i = 1, \dots, k\}$. Denote the stopping time $\tau_i := \inf\{t > \tau_{i-1}; X_t \in \Gamma \setminus \tilde{X}_{\tau_{i-1}}\}$ and $\tilde{X}_t := X_{\tau_{i-1}} \in \Gamma$ for $t \in [\tau_{i-1}, \tau_i)$ is defined by the sequence of $\tau_i, i = 0, 1, \dots$. This is the induced continuous time Markov chain

on Γ . The transition probability for \tilde{X} can be approximated by the large deviation principle for exit problems,

$$(C.1) \quad P_{i,i+1}^\varepsilon := \mathbb{P}\{\tilde{X} = i + 1 | \tilde{X} = i\} \approx ce^{-\frac{U_{i+\frac{1}{2}} - U_i}{\varepsilon}}.$$

Similarly, define $P_{i,i-1}^\varepsilon \propto e^{-\frac{U_{i-\frac{1}{2}} - U_i}{\varepsilon}}$. This defines an approximated Q -process with transition probability matrix (P_{ij}^ε) . Then the invariant distribution ν_i^ε , $i = 1, \dots, k$, satisfies

$$(C.2) \quad \nu_i^\varepsilon P_{i,i-1}^\varepsilon + \nu_i^\varepsilon P_{i,i+1}^\varepsilon = \nu_{i-1}^\varepsilon P_{i-1,i}^\varepsilon + \nu_{i+1}^\varepsilon P_{i+1,i}^\varepsilon.$$

One can directly verify the closed formula for ν_i^ε is given by

$$(C.3) \quad \nu_i^\varepsilon = \sum_{j=1}^k e^{-\frac{\tilde{h}_R(x_i; x_{j+1}) + \tilde{h}_L(x_i; x_j)}{\varepsilon}}.$$

Indeed, this formula is the principal left eigenvector $\nu^T Q = 0$ of a cyclic stochastic matrix

$$Q = \begin{pmatrix} -a_1 - b_1 & a_1 & & & b_1 \\ b_2 & -a_2 - b_2 & a_2 & & \\ & \ddots & \ddots & \ddots & \\ & & \ddots & \ddots & a_{k-1} \\ a_k & & & b_k & -a_k - b_k \end{pmatrix}, \quad \nu_i = \sum_{j=1}^k \prod_{\ell=i-1}^{i+j-1} b_\ell \prod_{m=i+1}^{i+j-1} a_m,$$

where $a_i = P_{i,i+1}^\varepsilon$ and $b_i = P_{i,i-1}^\varepsilon$ with k -periodic index.

Then as $\varepsilon \rightarrow 0$, by the Laplace principle for (C.3), we have

$$(C.4) \quad -\varepsilon \log \nu_i^\varepsilon \rightarrow W_i = \min_{j=1, \dots, k} (\tilde{h}_R(x_i; x_{j+1}) + \tilde{h}_L(x_i; x_j)), \quad i = 1, \dots, k.$$

This is the variational formula for boundary data W_i in (3.22).

Based on the invariant measure ν_i^ε , $i = 1, \dots, k$, for the induced Markov chain, one can recover the original invariant measure π_ε by the celebrated ergodic result by Khasminskii [Km60]. This, together with boundary data W_i , $i = 1, \dots, k$, can recover the Freidlin–Wentzell variational formula (3.28).

Acknowledgments. The authors would like to thank Jin Feng and Hung Tran for valuable suggestions.

REFERENCES

- [AK15] D. F. ANDERSON AND T. G. KURTZ, *Stochastic Analysis of Biochemical Systems*, Springer, New York, 2015.
- [Arn13] V. I. ARNOL'D, *Mathematical Methods of Classical Mechanics*, Grad. Texts in Math. 60, Springer, New York, 2013.
- [Aub83] S. AUBRY, *The twist map, the extended Frenkel-Kontorova model and the devil's staircase*, Phys. D, 7 (1983), pp. 240–258.
- [BD+97] M. BARDI AND I. CAPUZZO DOLCETTA, *Optimal Control and Viscosity Solutions of Hamilton-Jacobi-Bellman Equations*, Mod. Birkhäuser Class. 12, Birkhäuser/Springer, Basel, 1997.
- [BJ90] E. N. BARRON AND R. JENSEN, *Semicontinuous viscosity solutions for Hamilton-Jacobi equations with convex hamiltonians*, Comm. Partial Differential Equations, 15 (1990), pp. 293–309.

- [BS00] G. BARLES AND P. E. SOUGANIDIS, *On the large time behavior of solutions of Hamilton-Jacobi equations*, SIAM J. Math. Anal., 31 (2000), pp. 925–939.
- [CGMT15] F. CAGNETTI, D. GOMES, H. MITAKE, AND H. V. TRAN, *A new method for large time behavior of degenerate viscous Hamilton-Jacobi equations with convex Hamiltonians*, Ann. Inst. H. Poincaré Anal. Non Linéaire, 32 (2015), pp. 183–200.
- [CI99] G. CONTRERAS AND R. ITURRIAGA, *Global minimizers of autonomous lagrangians*, in Proceedings of the 22nd Brazilian Mathematics Colloquium, 1999.
- [CL83] M. G. CRANDALL AND P.-L. LIONS, *Viscosity solutions of Hamilton-Jacobi equations*, Trans. Amer. Math. Soc., 277 (1983), pp. 1–42.
- [CL84] M. G. CRANDALL AND P.-L. LIONS, *Two approximations of solutions of Hamilton-Jacobi equations*, Math. Comput., 43 (1984), pp. 1–19.
- [Con01] G. CONTRERAS, *Action potential and weak kam solutions*, Calc. Var. Partial Differential Equations, 13 (2001), pp. 427–458.
- [DFIZ16] A. DAVINI, A. FATHI, R. ITURRIAGA, AND M. ZAVIDOVIQUE, *Convergence of the solutions of the discounted Hamilton-Jacobi equation*, Invent. Math., 206 (2016), pp. 29–55.
- [E99] W. E, *Aubry-Mather theory and periodic solutions of the forced Burgers equation*, Comm. Pure Appl. Math., 52 (1999), pp. 811–828.
- [EGM84] P. EMBID, J. GOODMAN, AND A. MAJDA, *Multiple steady states for 1-D transonic flow*, SIAM J. Sci. Stat. Comput., 5 (1984), pp. 21–41.
- [ELVE19] W. E, T. LI, AND E. VANDEN-EIJNDEN, *Applied Stochastic Analysis*, Grad. Stud. Math. 199, AMS, Providence, RI, 2019.
- [Eva08] L. EVANS, *Weak KAM theory and partial differential equations*, in Calculus of Variations and Nonlinear Partial Differential Equations, Springer, New York, 2008, pp. 123–154.
- [Fat97] A. FATHI, *Théoreme KAM faible et théorie de mather sur les systemes lagrangiens*, C. R. Acad. Sci. Ser. I Math., 324 (1997), pp. 1043–1046.
- [Fat98] A. FATHI, *Sur la convergence du semi-groupe de Lax-Oleinik*, C. R. Acad. Sci. Ser. I Math., 327 (1998), pp. 267–270.
- [Fat08] A. FATHI, *Weak KAM Theorem in Lagrangian Dynamics Preliminary Version Number 10*, preprint, 2008, https://www.math.u-bordeaux.fr/~pthieull/Recherche/KamFaible/Publications/Fathi2008_01.pdf
- [FG12] A. FAGGIONATO AND D. GABRIELLI, *A representation formula for large deviations rate functionals of invariant measures on the one dimensional torus*, Ann. Inst. Henri Poincaré Probab. Stat., 48 (2012), pp. 212–234.
- [FR12] A. FIGALLI AND L. RIFFORD, *Aubry sets, Hamilton-Jacobi equations, and the Mañé conjecture*, Geom. Anal. Math. Relativity Nonlinear Partial Differential Equations, 599 (2012), pp. 83–104.
- [FRF09] A. FATHI, L. RIFFORD, AND A. FIGALLI, *On the Hausdorff dimension of the Mather quotient*, Comm. Pure Appl. Math., 62 (2009), pp. 445–500.
- [FW12] M. I. FREIDLIN AND A. D. WENTZELL, *Random Perturbations of Dynamical Systems*, Grundlehren Math. Wiss. 260, Springer, Berlin, 2012.
- [GL22] Y. GAO AND J.-G. LIU, *Revisit of macroscopic dynamics for some non-equilibrium chemical reactions from a hamiltonian viewpoint*, J. Stat. Phys., 189 (2022), pp. 1–57.
- [GL23] Y. GAO AND J.-G. LIU, *Large deviation principle and thermodynamic limit of chemical master equation via nonlinear semigroup*, Multiscale Model. Simul., to appear.
- [GLLL23] Y. GAO, T. LI, X. LI, AND J.-G. LIU, *Transition path theory for Langevin dynamics on manifold: Optimal control and data-driven solver*, Multiscale Model. Simul., 21 (2023), pp. 1–33.
- [Gom08] D. A. GOMES, *Generalized mather problem and selection principles for viscosity solutions and Mather measures*, Adv. Calc. Var., 1 (2008), pp. 291–307.
- [GQ12] H. GE AND H. QIAN, *Landscapes of non-gradient dynamics without detailed balance: Stable limit cycles and multiple attractors*, Chaos, 22 (2012), 023140.
- [II09] N. ICHIHARA AND H. ISHII, *Long-time behavior of solutions of Hamilton-Jacobi equations with convex and coercive hamiltonians*, Arch. Ration. Mech. Anal., 194 (2009), pp. 383–419.
- [IMT17] H. ISHII, H. MITAKE, AND H. V. TRAN, *The vanishing discount problem and viscosity mather measures. Part 1: The problem on a torus*, J. Math. Pures Appl., 108 (2017), pp. 125–149.
- [IS20] H. ISHII AND A. SICONOLFI, *The vanishing discount problem for Hamilton-Jacobi equations in the euclidean space*, Comm. Partial Differential Equations, 45 (2020), pp. 525–560.

- [Km60] R. Z. KHAS'MINSKII, *Ergodic properties of recurrent diffusion processes and stabilization of the solution to the Cauchy problem for parabolic equations*, Theory Probab. Appl., 5 (1960), pp. 179–196.
- [LPV86] P.-L. LIONS, G. PAPANICOLAOU, AND S. R. S. VARADHAN, *Homogenization of Hamilton-Jacobi Equations*, unpublished preprint, 1986.
- [Man92] R. MANÉ, *On the minimizing measures of lagrangian dynamical systems*, Nonlinearity, 5 (1992), pp. 623–638.
- [Man96] R. MANÉ, *Generic properties and problems of minimizing measures of lagrangian systems*, Nonlinearity, 9 (1996), pp. 273–310.
- [Mat82] J. N. MATHER, *Existence of quasi-periodic orbits for twist homeomorphisms of the annulus*, Topology, 21 (1982), pp. 457–467.
- [MT17] H. MITAKE AND H. V. TRAN, *Selection problems for a discount degenerate viscous Hamilton-Jacobi equation*, Adv. Math., 306 (2017), pp. 684–703.
- [NR99] G. NAMAH AND J.-M. ROQUEJOFFRE, *Remarks on the long time behaviour of the solutions of Hamilton-Jacobi equations*, Comm. Partial Differential Equations, 24 (1999), pp. 883–893.
- [Shu88] C.-W. SHU, *Total-variation-diminishing time discretizations*, SIAM J. Sci. Stat. Comput., 9 (1988), pp. 1073–1084.
- [SJ82] J. STEINHOFF AND A. JAMESON, *Multiple solutions of the transonic potential flow equation*, AIAA J., 20 (1982), pp. 1521–1525.
- [Sor15] A. SORRENTINO, *Action-minimizing Methods in Hamiltonian Dynamics (MN-50): An Introduction to Aubry-Mather Theory*, Princeton University Press, Princeton, NJ, 2015.
- [Tra21] H. V. TRAN, *Hamilton-Jacobi Equations: Theory and Applications*, Grad. Stud. Math. 213, AMS, Providence, RI, 2021.
- [Var66] S. R. S. VARADHAN, *Asymptotic probabilities and differential equations*, Comm. Pure Appl. Math., 19 (1966), pp. 261–286.
- [Var16] S. R. S. VARADHAN, *Large Deviations*, Courant Lect. Notes 27, AMS, Providence, RI, 2016.
- [VF69] A. D. VENTSEL' AND M. I. FREIDLIN, *The limiting behavior of the invariant measure under small perturbations of a dynamical system*, Dokl. Akad. Nauk, 188 (1969), pp. 13–16.
- [VF70] A. D. VENTSEL' AND M. I. FREIDLIN, *On small random perturbations of dynamical systems*, Russian Math. Surveys, 25 (1970), pp. 1–55.
- [WWY19] K. WANG, L. WANG, AND J. YAN, *Aubry-Mather theory for contact hamiltonian systems*, Comm. Math. Phys., 366 (2019), pp. 981–1023.
- [ZL16] P. ZHOU AND T. LI, *Construction of the landscape for multi-stable systems: Potential landscape, quasi-potential, A-type integral and beyond*, J. Chem. Phys., 144 (2016), 094109.